





COSMOSIL



COSMOSIL COSMOCORE

High Performance Liquid Chromatography

10th Edition





COSMOSIL Applications

COSMOSIL Application has more than 7,600 applications using COSMOSIL columns. Setting optimal HPLC experimental parameters is an important process that requires experience and time. COSMOSIL Applications provide you with sample analysis conditions for widely used ODS columns and our specialty columns.

- Over 7,600 applications
- Easy to search

Visit COSMOSIL top page at
<http://www.nacalai.co.jp/global/cosmosil/>

COSMOSIL HPLC Columns

General Info. of COSMOSIL/COSMOGEL
 COSMOSIL Columns List by Phase

- Standard Reversed Phase Columns
- Specialty Reversed Phase Columns
- Ultra-High Performance Columns
- Normal Phase Columns

Related Products

- Preparative Packing Materials
- Related Products

Hydrophilic Interaction Columns

Saccharide Separation Columns

Protein Separation Columns (Wide Pore Columns)

Fullerene Separation Columns

Carbon Nanotubes Separation Columns

COSMOSIL Applications
 Application Search
 Over 7,000 Data

Click

Sample Name contains (Keyword search) ▼

CAS number (ex:498-02-2)

Category (If no checkbox is clicked, the search will be performed in all categories.)

☐ Amino acids & derivatives ☐ Peptides & Proteins ☐ Nucleic acids & relative compounds

☐ Drugs & related compounds ☐ Antibiotics ☐ Vitamins

☐ Steroids ☐ Indoles ☐ Natural products

☐ Carbohydrates & derivatives ☐ Glyceride ☐ Oil

Column name (If no checkbox is clicked, the search will be performed in all columns.)

☐ C18-EB ☐ C18-MS-II ☐ C18-AR-II ☐ C18-PAQ

☐ COSMOCORE C18 ☐ Cholesterol ☐ PFP ☐ mNAP

☐ PYE ☐ NPE ☐ PBr ☐ CN-MS

☐ C8-MS ☐ C4-MS ☐ TMS-MS ☐ PE-MS

☐ SL-II ☐ HILIC ☐ Sugar-D ☐ NH2-MS

Particle Size ALL ▼

Application No. (ex:AP-1206)

Result/Page 20 ▼

Click

- Applications are searched by
1. Sample Category
 2. Sample Name
 3. CAS No.
 4. Column Name
 5. Particle Size

Search Result

COSMOSIL Application

Search condition (Application No:AP-1206)


[TOP]

Results 1 (1-1)

Data No.	Data Name	Particle Size	Column
AP-1206	Dichlorophenol	5	mNAP
2	Chlorophenol		576-24-9
2	Chlorophenol		120-63-2
2	Chlorophenol		583-78-8
2,6-Dichlorophenol			87-65-0
3,4-Dichlorophenol			95-77-2
3,5-Dichlorophenol			591-35-5


Click

COSMOSIL Application



INDEX

COSMOSIL / COSMOGEL Packing Material List	2	III. Preparative Packing Materials.....	60
Column Selection Guide	3	Normal and Reversed Phase Packing Materials	60
I. HPLC Columns.....	4	COSMOSIL C ₁₈ -OPN	61
1. Core-Shell Columns	4	COSMOSIL C ₁₈ -PREP	62
(1) COSMOCORE Series	4	Silica Gel (Spherical, Neutral)	63
COSMOCORE 2.6C ₁₈	6	Silica Gel (for Column Chromatography)	64
COSMOCORE 2.6Cholester	8	IV. Related Products.....	65
COSMOCORE 2.6PBr	10	1. Reagents for Mobile Phase Preparation	65
(2) Instrument Settings and Compatibility	12	Phosphate Buffer Solution (pH 2.5) (5X)	65
2. HPLC Columns	14	Stock Solutions for HPLC	65
(1) Reversed Phase Columns	14	Premixed Eluents for HPLC	65
C ₁₈ (ODS) Series	14	Additives	66
COSMOSIL C ₁₈ -MS-II	15	Ion-pair Reagents	66
COSMOSIL C ₁₈ -AR-II	17	2. Products for Sample Preparation	67
COSMOSIL C ₁₈ -PAQ	19	Cosmonice Filter	67
COSMOSIL C ₁₈ -EB	21	Cosmospin Filter	67
Reversed Phase Specialty Columns	22	Labeling Reagents	68
COSMOSIL Cholester	23	3. Column Care Products	69
COSMOSIL PBr	25	Cleaning Solution Kit	
COSMOSIL πNAP	26	for Reversed Phase HPLC Columns	69
COSMOSIL PYE	28	Storage Solution	
COSMOSIL NPE	29	for Reversed Phase HPLC Columns	70
Other Reversed Phase Columns	30	4. COSMOSIL HPLC Accessories	71
COSMOSIL PFP	31		
COSMOSIL CN-MS	32		
COSMOSIL C ₂₂ -AR-II, C ₈ -MS, C ₄ -MS,			
TMS-MS, PE-MS	33		
(2) Normal Phase Columns	34		
COSMOSIL SL-II	34		
(3) Hydrophilic Interaction Columns	35		
COSMOSIL HILIC	35		
(4) Mono- and Oligosaccharide Analysis Columns	37		
COSMOSIL Sugar-D	38		
COSMOSIL NH ₂ -MS	39		
(5) Protein Separation Columns	40		
Reversed Phase Columns	40		
COSMOSIL Protein-R	40		
COSMOSIL C ₁₈ -AR-300, C ₈ -AR-300,			
C ₄ -AR-300, Ph-AR-300	41		
Gel Filtration Columns (Aqueous)	43		
COSMOSIL Diol-120-II, Diol-300-II, Diol-1000-II	43		
Ion Exchange Columns	45		
COSMOGEL IEX Series	45		
Hydrophobic Interaction Columns	47		
COSMOSIL HIC	47		
(6) Columns for Fullerene Separation	48		
COSMOSIL Buckyprep	49		
COSMOSIL Buckyprep-D	50		
COSMOSIL Buckyprep-M	51		
COSMOSIL PBB	52		
COSMOSIL NPE	52		
COSMOSIL PYE	53		
(7) Columns for Soluble Carbon Nanotube Separation	54		
COSMOSIL CNT-300, CNT-1000, CNT-2000	54		
II. SFC Columns.....	55		
COSMOSIL HP (3-Hydroxyphenyl)	56		
COSMOSIL PY (Pyridinyl)	57		
COSMOSIL Quinoline	58		
COSMOSIL Cholester	59		
COSMOSIL PBr	59		



COSMOSIL / COSMOGEL Packing Material List

Core-Shell Columns

Sample	Separation Mode	Packing Material	Bonded Phase	Bonding Type	Average Particle Size (μm)	Average Pore Size (Å)	Carbon Content (%)	Special Features and Applications	USP Category	Page
Organic compounds (low M.W.)	Reversed phase	C ₁₈	Octadecyl group	Polymeric	2.6	90	7	Multi-purpose C ₁₈ column	L1	5, 6
		Cholester	Cholesteryl group	Mono-meric			-	Usable under the same condition as C ₁₈ . Unique rigid cholesteryl structure improves separation.	L101	5, 8
		PBr ^{NEW}	Pentabromobenzyl group					Separate hydrophilic compounds under reversed-phase conditions.	-	5,10

HPLC Columns


Sample	Separation Mode	Packing Material	Bonded Phase	Bonding Type	Average Particle Size (μm)	Average Pore Size (Å)	Carbon Content (%)	Special Features and Applications	USP Category	Page
Organic compounds (low M.W.)	Reversed phase	C ₁₈ -MS-II	Octadecyl group	Mono-meric	2.5	130	18	Multi-purpose C ₁₈ column	L1	14, 15
				3, 5, 15	120	16				
		C ₁₈ -AR-II		Polymeric	3, 5, 15	120	17	Features strong acid resistance, good for acidic compounds and peptides.		14, 17
		C ₁₈ -PAQ			5, 15		11	Good for hydrophilic compounds, and stable performance under 100% aqueous conditions.		14, 19
		C ₁₈ -EB	Mono-meric	3	14.5	Good for basic compounds	14, 21			
		Cholester	Cholesteryl group	Mono-meric	2.5	130	21	Usable under the same conditions as C ₁₈ . Unique rigid cholesteryl structure improves separation.	L101	22, 23
			5		120	20				
		PBr	Pentabromobenzyl group		5	120	8	Separate hydrophilic compounds under reversed-phase conditions.	-	22, 25
		πNAP	Naphthylethyl group		2.5	130	14	Stronger π-π interaction than phenyl column.		22, 26
			5		120	11		22, 28		
		PYE	Pyrenylethyl group		5	120	18	The most powerful π-π interaction		
		NPE	Nitrophenylethyl group	9			Separation utilizing dipole-dipole interaction.			
		PFP	Pentafluorophenyl group	10			Separation utilizing weak dipole-dipole interaction.	L43	30, 31	
		CN-MS	Cyanopropyl group	7			Enables separation of different hydrophobic samples without using gradients.	L10	30, 32	
		C ₂₂ -AR-II	Docosyl group	Polymeric	19	Alkyl chain columns, excluding C ₁₈ column	-	30, 33		
		C ₈ -MS	Octyl group	10	L7					
	C ₄ -MS	Butyl group	7	L26						
	TMS-MS	Trimethyl group	5	L13						
	PE-MS	Phenylethyl group		10	π-π interaction	L11				
	Normal phase	SL-II	--	-	3, 5, 15	120	-	Suitable for preparative separation.	L3	34
Hydrophilic interaction	HILIC	Triazole	-	2.5	130	-	Retains highly polar compounds that would not be retained in a C ₁₈ olumn.	L104	35	
				5	120					
Mono- and Oligo-saccharides	Hydrophilic interaction	Sugar-D	Secondary/Tertiary amine	-	5	-	-	A novel stationary phase for mono- and oligosaccharides.	-	37, 38
		NH ₂ -MS	Aminopropyl group	Polymeric		120	4	Primary amino bonded column	L8	37, 39
Proteins	Reversed phase	Protein-R	Octadecyl group	Polymeric	5	300	-	Wide-pore column with the advantages of both C ₁₈ and C ₄	L1	40
		C ₁₈ -AR-300					12	Wide pore type	L1	41
		C ₈ -AR-300	7				L7			
		C ₄ -AR-300	6				L26			
		Ph-AR-300	7				L11			
	Gel permeation	Diol-120-II	Diol group	-	5	120	-	Silica-based gel filtration column Sample MW (Protein) 5,000-100,000 Da	L20	43
		Diol-300-II				300		Silica-based gel filtration column Sample MW (Protein) 10,000-700,000 Da		
		Diol-1000-II				1000		Silica-based gel filtration column Sample MW (Water Soluble Polymer) 50,000-500,000 Da		
	Ion-exchange	IEX Type Q	Trimethylaminopropyl type	-	5	1000	-	Anion-exchange type (purification)		45
		IEX Type Q-N				-		Anion-exchange type (ultra-fast analysis, precise analysis)		
		IEX Type S	Sulfopropyl type			1000		Cation-exchange type (purification)		
		IEX Type S-N				-		Cation-exchange type (ultra-fast analysis, precise analysis)		
		IEX Type M	Trimethylaminopropyl type /Sulfopropyl type			1000		Amphoteric ion-exchange type (purification)		
		IEX Type M-N				-		Amphoteric ion-exchange type (precise analysis)		
	Hydrophobic interaction	HIC	--	-	5	300	-	Little loss in enzyme activity and the tertiary structure of proteins		47

Sample	Separation Mode	Packing Material	Bonded Phase	Bonding Type	Average Particle Size (μm)	Average Pore Size (Å)	Carbon Content (%)	Special Features and Applications	USP Category	Page
Fullerenes	--	Buckyprep	Pyrenylpropyl group	Mono-meric	5	120	17	Standard column for fullerene separation.		48, 49
		Buckyprep-D	Nitro-carbazoyl group				-	Good for derivatized fullerenes		48, 50
		Buckyprep-M	Phenothiazinyl group				13	Good for metallofullerenes		48, 51
		PBB	Pentabromobenzyl group				8	Good for preparative separation of C ₆₀ or C ₇₀ .		48, 52
		NPE	Nitrophenylethyl group				9	Separation of derivatized fullerenes		48, 53
		PYE	Pyrenylethyl group				18	Separation of fullerenes		48, 52
Carbon nanotubes	Gel permeation	CNT-300	Hydrophilic group (neutral)	-	5	300	-	Separation of soluble carbon nanotubes.		54
		CNT-1000				1000				
		CNT-2000				2000				

SFC Columns

Sample	Separation Mode	Packing Material	Bonded Phase	Bonding Type	Average Particle Size (μm)	Average Pore Size (Å)	Carbon Content (%)	Special Features and Applications	USP Category	Page
-	SFC	HP ^{NEW}	3-Hydroxyphenyl group	Polymeric	3, 5	120	-	Good for hydrophilic compounds. Stronger retention for basic compounds than PY	-	55, 56
		PY ^{NEW}	Pyridinyl group					Similar separation properties as 2-Ethylpyridine, with stronger retention.		55, 57
		Quinoline ^{NEW}	Quinoline group	Mono-meric	2.5, 5	130	21	Alternate selectivity to HP and PY.	L101	55, 58
		Cholester	Cholesteryl group		2.5	130		Usable under the same conditions as C ₁₈ . Unique rigid cholesteryl structure improves separation.		55, 59
		PBr	Pentabromobenzyl group		5	120		Separate hydrophilic compounds under reversed-phase conditions.		55, 59

Column Selection Guide




I. HPLC Columns

1. Core-Shell Columns

(1) COSMOCORE Series

About Core-Shell Particles


Core-shell particles consist of a nonporous core inside a porous shell. By using these core-shell particles, one can achieve sharper peaks compared to fully porous silica gel particles of the same diameter.




Schematic diagram of a silica particle

Uniform Particle Size Distribution Compared to 1.7 μm Particles

Compared to fully porous particles, core-shell particles have a more uniform particle diameter; therefore, core-shell particles can be packed in the column more uniformly to minimize sample diffusion.




COSMOCORE 2.6C18 (200x)




Fully porous 1.7 μm particles (200x)

Mass Transfer Equivalent to Fully Porous sub-2 μm Particles

Mass transfer refers to the time it takes for a sample molecule to enter and leave a particle. In general, lower mass transfer time corresponds to less diffusion and sharper peaks. Even though COSMOCORE 2.6C18 has a larger particle diameter than fully porous sub-2 μm particles, the mass transfer characteristics are similar.





COSMOCORE 2.6C18



Fully porous sub-2 μm particle

Reduced Back Pressure and Faster Analyses

COSMOCORE 2.6C18 delivers performance equivalent to sub-2 μm particles at faster flow rate and analysis time while maintaining a lower back pressure. COSMOCORE can also be used in longer column size to gain additional resolution.




Column size: 2.1 mm I.D. x 50 mm
Mobile phase: Acetonitrile / Water = 70 / 30

Temperature: 40°C
Sample: Amylbenzene



Same performance and lower back pressure compared to sub-2 µm particles

Reduced Back Pressure

COSMOCORE 2.6C₁₈ maintains the same performance as sub-2 µm particles with half the back pressure.



Specifications


Packing Material	2.6C ₁₈	2.6Cholester	2.6PBr
Silica Gel	Core-shell type silica gel		
Average Particle Size	2.6 μm		
Average Core Diameter	1.6 μm		
Average Pore Size	approx. 90 Å		
Specific Surface Area	approx. 150 m ² /g		
Bonded Phase Structure			
Bonded Phase	Octadecyl group	Cholesteryl group	Pentabromobenzyl Group
Main interaction	Hydrophobic interaction	Hydrophobic interaction Molecular shape selectivity	Hydrophobic interaction Dispersion force
End-Capping Treatment	Near-perfect treatment		
Usable pH Range	1.5 - 10	2 - 7.5	
Maximum Pressure	60 MPa		

COSMOCORE 2.6C₁₈

- Ultra-high performance LC results with conventional HPLC equipment
- Same number of theoretical plates as sub-2 µm columns with half the back pressure
- Increased loading capacity
- Excellent pH stability (1.5-10)


Excellent pH Stability

Under accelerated pH 10.4, 40°C stability test, COSMOCORE C₁₈ shows superior stability compared with other core-shell C₁₈ phases.



Column size: 2.1 mm x 100 mm
Sample: Caffeine (0.05 mg/ml)
Flow rate: 0.4 ml/min.

Mobile phase: 0.35% Ammonium hydroxide/ acetonitrile = 90/10 (pH 10.4)
Injection volume: 1 µl
Temperature: 40 °C





Sharp Peaks with Many Types of Compounds

COSMOCORE 2.6C₁₈ features a special end capping treatment that effectively shields residual silanol groups, yielding sharp peaks for basic compounds and metal coordination complexes.

Basic Compounds



• Amitriptyline

Column size: 2.1 mm I.D. - 50 mm
Mobile phase: Acetonitrile / 20mmol/l Phosphoric Buffer(pH7.0) = 60 / 40
Temperature: 40°C
Flow rate: 0.4 ml/min
Detection: UV254nm
Sample: 1; Amitriptyline (0.2 mg/ml)
2; Amylbenzene I.S. (2.0 mg/ml)
Inj. Vol.: 0.5 µl





• Haloperidol

Column size: 2.1 mm I.D. - 50 mm
Mobile phase: A; Acetonitrile/ 20mmol/l Phosphate buffer(pH7.0) = 10/90
B; Acetonitrile/ 20mmol/l Phosphate buffer(pH7.0) = 50/50
B conc. 0→100%(0→3min), 100%(3-5min)
Flow rate: 0.4 ml/min
Temperature: 40°C
Detection: UV220nm
Sample: 1; 4-(4-Chlorophenyl)-4-hydroxypiperidine (0.1 mg/ml)
2; Haloperidol (0.1 mg/ml)
Inj. Vol: 0.5 µl





• Procainamide

Column size: 2.1 mm I.D. - 50 mm
Mobile phase: Acetonitrile/ 20mmol/l Phosphate buffer(pH7.0) = 10/90
Flow rate: 0.4 ml/min
Temperature: 40°C
Detection: UV270nm
Sample: 1; Procainamide (0.05 mg/ml)
2; N-Acetylprocainamide (0.05 mg/ml)
Inj. Vol: 0.5 µl




• Caffeine

Column size: 2.1 mm I.D. - 100 mm
Mobile phase: Acetonitrile/ H₂O = 10/90
Flow rate: 0.4 ml/min
Temperature: 40°C
Detection: UV275nm
Sample: 1; Theobromine (0.05 mg/ml)
2; Theophylline (0.05 mg/ml)
3; Caffeine (0.05 mg/ml)
Inj. Vol.: 1.0 µl




Metal Coordination Complexes

• Oxine-Copper



Low Bleed-Suitable for LC-MS

COSMOCORE 2.6C₁₈ has low column bleed and consequently low MS noise level.



Ordering Information

• Analytical Columns (Particle Size: 2.6 µm)

COSMOCORE 2.6C₁₈ Packed Columns

Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
2.1 x 30	12632-31	3.0 x 30	12611-01	4.6 x 30	12601-31
2.1 x 50	12631-41	3.0 x 50	12609-51	4.6 x 50	12600-41
2.1 x 75	12630-51	3.0 x 75	12608-61	4.6 x 75	12599-91
2.1 x 100	12614-71	3.0 x 100	12607-71	4.6 x 100	12598-01
2.1 x 150	12612-91	3.0 x 150	12602-21	4.6 x 150	12597-11
				4.6 x 250	12596-21

COSMOCORE's connector is the same type as Waters UPLC® columns.


For UHPLC-compatible prefilters, refer to page 13.

COSMOCORE 2.6Cholester

- Cholesterol-bonded reversed-phase core-shell column
- Usable under the same conditions as C₁₈ columns
- Better selectivity for cis-trans isomers, polyphenols, and natural products


Comparison with C₁₈

COSMOCORE 2.6Cholester offers improved separation for cis-trans isomers than C₁₈ under typical reversed-phase mobile phase.




Separation Properties

COSMOCORE 2.6Cholester has about the same hydrophobicity as C₁₈. It is not necessary to change the analytical conditions when replacing C₁₈ Columns with COSMOCORE 2.6Cholester.



Molecular Shape Selectivity

COSMOCORE 2.6Cholester has excellent shape selectivity due to its structural rigidity. COSMOCORE 2.6Cholester retains planar triphenylene longer than non planar *o*-terphenyl.



Applications


• Vitamin D metabolites

COSMOCORE Application Data

Column: COSMOCORE 2.6Cholester
 Column size: 2.1mm I.D.-150mm
 Mobile phase: A; 0.1% Formic acid-H₂O
 B; Methanol
 B conc. 75→82% (0→11min), 95% (11→25min)
 Flow rate: 0.3 ml/min
 Temperature: 50°C
 Detection: UV265nm

Sample:

- 1; 25-Hydroxyvitamin D₃
- 2; 3-*epi*-25-Hydroxyvitamin D₃
- 3; 25-Hydroxyvitamin D₂
- 4; 3-*epi*-25-Hydroxyvitamin D₂
- 5; Vitamin D₂ [Calciferol]
- 6; Vitamin D₃



Data courtesy of a customer

AP-1514


• Catechins

COSMOCORE Application Data

Column: COSMOCORE 2.6Cholester
 Column size: 2.1mm I.D.-50mm
 Mobile phase: A; Acetonitrile/ 0.1% H₃PO₄ = 10/90
 B; Acetonitrile/ 0.1% H₃PO₄ = 40/60
 B conc. 0→100% 3min Linear gradient
 Flow rate: 0.6 ml/min
 Temperature: 30°C
 Detection: UV280nm

Sample:

- 1; (-)-Gallocatechin[(-)-GC]
- 2; Caffeine
- 3; (-)-Epigallocatechin[(-)-EGC]
- 4; (-)-Catechin[(-)-C]
- 5; (-)-Epicatechin[(-)-EC]
- 6; (-)-Epigallocatechin Gallate[(-)-EGCg]
- 7; (-)-Gallocatechin Gallate[(-)-GCg]
- 8; (-)-Epicatechin Gallate[(-)-ECg]
- 9; (-)-Catechin Gallate[(-)-Cg]



Data courtesy of a customer

Ordering Information

• Analytical Columns (Particle Size: 2.6 µm)

COSMOCORE 2.6Cholester Packed Column

Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
2.1 x 30	12858-91	3.0 x 30	12863-11	4.6 x 30	12869-51
2.1 x 50	12859-81	3.0 x 50	12864-01	4.6 x 50	12870-11
2.1 x 75	12860-41	3.0 x 75	12866-81	4.6 x 75	12871-01
2.1 x 100	12861-31	3.0 x 100	12867-71	4.6 x 100	12872-91
2.1 x 150	12862-21	3.0 x 150	12868-61	4.6 x 150	12873-81
				4.6 x 250	12875-61

COSMOCORE's connector is the same type as Waters UPLC® columns.
 For UHPLC-compatible prefilters, refer to page 13.

COSMOCORE 2.6PBr

- Separate hydrophilic compounds under reversed-phase conditions
- Retain hydrophilic compounds longer than C₁₈
- Greater sample loading capacity than HILIC
- High performance similar to sub-2 µm particles with lower back pressure


< Suitable Samples >

- Hydrophilic compounds
- Nucleic acids and derivatives
- Surfactants
- Glycosides
- Peptides


Separation of Hydrophilic Compounds (low retention on C₁₈)

COSMOCORE 2.6PBr retains hydrophilic compounds stronger than C₁₈ columns under the same reversed-phase conditions.


• Nucleic Acid Metabolites



• Malonyl CoA, CoA, Acetyl CoA




• Water-Soluble Vitamins




Separation of Hydrophilic Compounds (compounds with similar hydrophobicity)

COSMOCORE 2.6PBr can separate compounds with similar hydrophobicity, utilizing several kinds of molecular interactions, including dispersion force generated by the bromine atoms.


• ATP, ADP, AMP



• Vitamin B3 (Nicotinic Acid, Nicotinamide)




• Thiamine Pyrophosphate, Thiamine




Separation of Hydrophilic Compounds (Glycosides)

Glycosides with identical aglycones but different glycosyl groups can also be separated.

• UDP Glycosides



• Arbutin and Hydroquinone



Ordering Information

• Analytical Columns (Particle Size: 2.6 µm)

COSMOCORE 2.6PBr Packed Column

Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
2.1 x 30	13692-21	3.0 x 30	13698-61	4.6 x 30	13705-51
2.1 x 50	13693-11	3.0 x 50	13699-51	4.6 x 50	13712-51
2.1 x 75	13694-01	3.0 x 75	13700-01	4.6 x 75	13714-31
2.1 x 100	13695-91	3.0 x 100	13701-91	4.6 x 100	13715-21
2.1 x 150	13697-71	3.0 x 150	13703-71	4.6 x 150	13719-81
				4.6 x 250	13734-71

COSMOCORE's connector is the same type as Waters UPLC® columns.

For UHPLC-compatible prefilters, refer to page 13.

(2) Instrument Settings and Compatibility

When using with a conventional (non-UHPLC) instrument


COSMOCORE columns are designed for use with UHPLC instruments. In addition, due to their low backpressure, they can be used with conventional instruments. However, it is necessary to change the following settings.

Detector Response Time

Because UHPLC analyses are done at high flow rates, a slow response time can adversely affect peak shape. We recommend setting the response time to 0.1 sec or less.

COSMOCORE Application Data

Column: COSMOCORE 2.6C₁₈
 Column size:
 Mobile phase: Acetonitrile/ Water = 70 /30
 Flow rate: 3.0mmI.D.-75mm 1.0 ml/min
 2.1mmI.D.-50mm 0.4ml/min
 Temperature: 40°C
 UV254nm
 Sample: 1; Benzene (2.0mg/ml)
 2; Toluene (2.0mg/ml)
 3; Ethylbenzene (2.0mg/ml)
 4; Propylbenzene (2.0mg/ml)
 5; Butylbenzene (2.0mg/ml)
 6; Amylbenzene (2.0mg/ml)
 Inj. Vol.: 3.0mmI.D.-75mm 1.0µl
 2.1mmI.D.-50mm 0.5µl




Other Instrument Parameters

UHPLC is more vulnerable to the effects of dead volume than conventional chromatography. When using a 2.1 mm I.D. column, please use a semi-micro detector cell, injector, and piping (0.1mm).

COSMOCORE Application Data

Column: COSMOCORE 2.6C₁₈
 Column size:
 Mobile phase: Acetonitrile/ Water = 70/30
 Flow rate: 3.0mmI.D.-75mm 1.0 ml/min
 2.1mmI.D.-50mm 0.4ml/min
 Temperature: 40°C
 Detection: UV254nm
 Sample: 1; Benzene (2.0mg/ml)
 2; Toluene (2.0mg/ml)
 3; Ethylbenzene (2.0mg/ml)
 4; Propylbenzene (2.0mg/ml)
 5; Butylbenzene (2.0mg/ml)
 6; Amylbenzene (2.0mg/ml)
 Inj. Vol.: 3.0mmI.D.-75mm 1.0µl
 2.1mmI.D.-50mm 0.5µl



Fittings and Adapters



COSMOCORE columns use the same connectors as Waters UPLC® (UHPLC) columns. This is different from our conventional COSMOSIL columns, which use the conventional Waters HPLC-compatible connectors.

(UPLC® is a registered trademark of Waters Corporation.)

1. Differences between end fitting

Connection Type		Column	
		HPLC(COSMOSIL)	UHPLC (COSMOCORE)
Instrument	HPLC	No adapter required	Adapter required
	UHPLC	Adapter required	No adapter required

HPLC: Conventional Waters-compatible connector
 UHPLC: Waters UPLC®-compatible connector


	HPLC	UHPLC
Connector Shape	 approx. 3.3 mm	 approx. 2.3 mm

The length of tubing that extends from the ferrule differs from HPLC to UHPLC.

2. COSMOCORE-compatible fittings

1) UHPLC instrument fittings

No adapter needed; just connect as-is.




2) HPLC instrument fittings

An adapter or movable (high-pressure) fitting is required to connect the fittings to the column. See the examples for different fittings below.


SUS ferrules (HPLC) fixed on the tubing

The column can be connected by using an adapter (SUS union + tubing fixed to UHPLC length).



PEEK fittings

PEEK fittings do not fix the length of tubing at the end, so they can be used with both types of column. However, please be cautious of their pressure tolerance.



Ordering Information

• Adapter List

Product Name	Description	Product Number	PKG Size
Low & Zero Dead Volume Union	Material: SUS Bore diameter: 0.35 mm	P0402	1 PKG
COSMOSIL Column Connecting Tube (0.1 mm I.D.)	I.D.: 0.1 mm	12570-41	1 PKG
COSMOSIL Column Connecting Tube (0.25 mm I.D.)	I.D.: 0.25 mm	37843-69	1 PKG

• UHPLC-compatible prefilter



Product Name	In	Out	Contents	Product Number	PKG Size
U-Fil UHPLC-compatible prefilter	UHPLC	UHPLC	Filter: 0.5 μ m	12571-31	1 SET
	HPLC	UHPLC	Tubing connecting diameter: 1/16	12572-21	1 SET
U-Fil replacement filter	-	-	Filter: 0.5 μ m Material: SUS316L	15767-91	5 units / PKG

2. HPLC Columns

(1) Reversed Phase Columns

C₁₈ (ODS) Series


Specifications

Packing Material	C18-MS-II	C18-AR-II	C18-PAQ	C18-EB
Silica Gel	High purity porous spherical silica			
Average Particle Size	2.5, 3 , 5, 15 μm	3, 5, 15 μm	5, 15 μm	3 μm
Average Pore Size	approx. 120 Å			
Specific Surface Area	approx. 300 m ² /g			
Bonded Phase Structure				
Bonded Phase	Octadecyl group			
Bonding Type	Monomeric	Polymeric		Monomeric
Main Interaction	Hydrophobic interaction			
End-Capping Treatment	Near-perfect treatment			
Carbon Content	approx. 16%	approx. 17%	approx. 11%	approx. 14.5%
Usable pH Range	2~10*	1.5~7.5*	2~7.5	2-10*
Features	•Multi-purpose C18 Column	•Features strong acid resistance. •Good for acidic compounds and peptides.	•Good for hydrophilic compounds. •Stable performance under 100% aqueous conditions.	•Good for basic compounds

*Optimal pH range of silica-based columns is between 2 and 7.5. Extreme pH may significantly decrease column lifetime.

Difference in Separation Properties (5 μm)

COSMOSIL 5C₁₈-AR-II retains planar compounds (such as triphenylene) longer compared to COSMOSIL 5C₁₈-MS-II. COSMOSIL 5C₁₈-PAQ has shorter retention time, and retains polar compounds (Such as valerophenone, *n*-butyl benzoate) longer.




COSMOSIL C₁₈-MS-II

- First-choice column of our ODS series
- Multi-purpose C₁₈ column
- High reproducibility
- A wide range of applications

< Suitable Samples >

- Low-M.W. Compounds



Separation Property

The COSMOSIL 5C₁₈-MS-II is a well-balanced column with better basic performance, such as sharper peaks for basic compounds and chelating compounds, strong hydrophobic interaction, low analytical pressure, and high theoretical plate number. COSMOSIL 5C₁₈-MS-II is the first-choice column for reversed-phase chromatography.






Table. Comparison of hydrophobic interaction, analytical pressure, and theoretical plate number

Column	Hydrophobic Interaction α (Toluene/Benzene)	Pressure (MPa)	Theoretical Plate Number (Toluene)
COSMOSIL 5C ₁₈ -MS-II	1.96	8.3	14300
Company A C ₁₈	1.99	13.0	16800
Company B C ₁₈	1.94	8.0	14000
Company C C ₁₈	1.69	11.2	5600
Company D C ₁₈	1.84	10.5	14200

Analysis of Basic Compounds and Metal Coordination Compounds

The COSMOSIL 5C₁₈-MS-II column, taking advantage of a new end-capping treatment, can replace the original COSMOSIL C₁₈ (ODS) column. A new end-capping treatment with polar groups for shielding effect has significantly improved peak shape for basic compounds. Ultra pure silica gel with low trace-metal content is used for COSMOSIL columns; thus the columns provide excellent peak shapes for chelating compounds.




High Reproducibility

The strict quality control system of Nacalai Tesque ensures the quality of the silica gel and bonding and end capping process, reducing variation between lots. We support customers with an individual Inspection Report which accompanies each and every COSMOSIL, COSMOCORE and COSMOGEL packed column (except guard columns) and an additional Certificate of Analysis for the COSMOSIL 5C₁₈-MS-II (4.6 mm I.D. x 150 mm and 4.6 mm I.D. x 250 mm).


A Wide Range of Applications

A wide selection of applications, e.g. drug molecules, is available to achieve appropriate separation parameters for target samples.

• Parasympatholytic Agents



• Analgesic Antipyretic Drugs



Ordering Information

• Analytical / Preparative Columns (Particle Size: 5 µm)

COSMOSIL 5C₁₈-MS-II Packed Column

Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
1.0 x 50	02824-31	4.6 x 100	38018-91
1.0 x 150	02896-01	4.6 x 150*	38019-81
2.0 x 30	05876-71	4.6 x 150 3 lots set	09397-73
2.0 x 50	04355-21	4.6 x 250*	38020-41
2.0 x 100	05597-31	6.0 x 150	38021-31
2.0 x 150	38025-91	6.0 x 250	38022-21
2.0 x 250	05761-61	10 x 50	05789-21
3.0 x 100	05458-51	10 x 150	34355-91
3.0 x 150	34245-31	10 x 250	38023-11
3.0 x 250	34254-11	20 x 150	05091-41
4.6 x 30	34341-61	20 x 250	38024-01
4.6 x 50	38017-01	28 x 250	05760-71

COSMOSIL 5C₁₈-MS-II Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	38014-31
4.6 x 10 Cartridge**	38015-89
10 x 20	38016-11
20 x 20	05790-81
20 x 50	34371-71
28 x 50	34347-01

* Validated columns

** 2 cartridges included. Guard cartridge holder required; refer to page 71.

• Preparative Columns (Particle Size: 15 µm)

COSMOSIL 15C₁₈-MS-II Packed Column

Column Size I.D. x Length (mm)	Product Number
28 x 250	34525-61
50 x 250	05886-41
50 x 500	34531-71

COSMOSIL 15C₁₈-MS-II Guard Column

Column Size I.D. x Length (mm)	Product Number
28 x 50	05885-51
50 x 50	34527-41

• Fast LC Column (Particle Size: 3 µm)

COSMOSIL 3C₁₈-MS-II Packed Column

Column Size I.D. x Length (mm)	Product Number
2.0 x 50	05514-01
4.6 x 10	38065-71
4.6 x 50	38066-61
4.6 x 100	38067-51

• Analytical Columns (Particle Size: 2.5 µm)

COSMOSIL 2.5C₁₈-MS-II Packed Column


Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
2.0 x 50	08994-31	3.0 x 50	08997-01
2.0 x 75	08995-21	3.0 x 75	08998-91
2.0 x 100	08996-11	3.0 x 100	08999-81

COSMOSIL C₁₈-AR-II

- Features strong acid resistance


< Suitable Samples >

- Peptides, acidic compounds, etc.



Acid Resistance

The COSMOSIL 5C₁₈-AR-II packed column features a polymeric type of C₁₈ reversed phase material. The acidic resistance of COSMOSIL 5C₁₈-AR-II is much improved compared with commercially available monomeric type octadecyl stationary phases. It retains high performance even with acidic mobile phases commonly used to separate acidic compounds and peptides.



Decomposition test in 0.1% Trifluoroacetic acid solution at 60°C.
Capacity factor (*k'*) = Naphthalene,
Mobile phase: Methanol / H₂O=70/30


Applications

• Peptides

COSMOSIL Application Data

Column: 5C₁₈-AR-II
Column size: 4.6mm I.D.-150mm
Mobile phase: A; 0.05%TFA-H₂O
B; 0.05%TFA-Acetonitrile
B conc. 10→40% 20min Linear gradient
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV220nm

Sample: 1; Oxytocin (0.8μg)
2; Angiotensin II(Human) (0.8μg)
3; Angiotensin I(Human) (0.8μg)
4; Substance P (0.8μg)



NACALAI TESQUE, INC
AP-0351

• Organic Acids

COSMOSIL Application Data

Column: 5C₁₈-AR-II
Column size: 4.6mm I.D.-150mm
Mobile phase: Methanol/ 20mmol/l Phosphoric Acid
= 20/80
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV254nm, 0.5AUFS

Sample: 1; Gallic Acid (0.63μg)
2; Protocatechuic Acid (0.63μg)
3; Gentisic Acid (0.63μg)
4; Phthalic Acid (0.63μg)




NACALAI TESQUE, INC
AP-0159

• Salicylic Acid Esters

COSMOSIL Application Data

Column: 5C₁₈-AR-II
Column size: 4.6mm I.D.-150mm
Mobile phase: Acetonitrile/ H₂O = 70/30
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV254nm, 0.2AUFS

Sample: 1; Methyl Salicylate (2.3μg)
2; Ethyl Salicylate (2.6μg)
3; *n*-Propyl Salicylate (2.3μg)
4; *n*-Butyl Salicylate (2.6μg)




NACALAI TESQUE, INC
AP-0165

• Parabens

COSMOSIL Application Data

Column: 5C₁₈-AR-II
Column size: 4.6mm I.D.-150mm
Mobile phase: Acetonitrile/ H₂O = 50/50
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV254nm, 0.12AUFS


Sample: 1; Methyl *p*-Hydroxybenzoate (0.125μg)
2; Ethyl *p*-Hydroxybenzoate (0.125μg)
3; *n*-Propyl *p*-Hydroxybenzoate (0.125μg)
4; Butyl *p*-Hydroxybenzoate (0.125μg)



NACALAI TESQUE, INC
AP-0099

LC/MS Applications

- Identification of herbal medicine constituents by LC/MS



Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 µm)

COSMOSIL 5C₁₈-AR-II Packed Column

Column Size I.D. x Length (mm)	Product Number
1.0 x 50	02955-21
1.0 x 150	02951-61
2.0 x 30	05098-71
2.0 x 50	34400-81
2.0 x 100	34469-11
2.0 x 150	37992-51
2.0 x 250	05272-71
3.0 x 100	05791-71
3.0 x 150	38028-61
3.0 x 250	38029-51
4.6 x 30	05877-61
4.6 x 50	38142-51

COSMOSIL 5C₁₈-AR-II Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	38141-61
4.6 x 10 Cartridge**	38008-89
10 x 20	38148-91
20 x 20	34458-51
20 x 50	34479-81
28 x 50	34363-81

* Validated columns

** 2 cartridges included. Guard cartridge holder required; refer to page 71.

- Preparative Columns (Particle Size: 15 µm)

COSMOSIL 15C₁₈-AR-II Packed Column

Column Size I.D. x Length (mm)	Product Number
28 x 250	37978-51
50 x 250	38058-71
50 x 500	05884-61

COSMOSIL 15C₁₈-AR-II Guard Column

Column Size I.D. x Length (mm)	Product Number
28 x 50	38030-11
50 x 50	38057-81

- Fast LC Column (Particle Size: 3 µm)

COSMOSIL 3C₁₈-AR-II Packed Column


Column Size I.D. x Length (mm)	Product Number
2.0 x 50	05478-91
4.6 x 10	38068-41
4.6 x 50	38069-31
4.6 x 100	38070-91

COSMOSIL C₁₈-PAQ

- Compatible with 100% water based mobile phase


< Suitable Samples >

- Hydrophilic compounds
- Organic acids, nucleic acid bases, etc.



Stable Performance


Stable performance under 100% aqueous conditions




The figure shows the change of retention time for thymine with 100% aqueous mobile phase (20 mmol/l phosphate buffer, pH 7). The sample was analyzed 4 times (1 hour). Flow of mobile phase was then stopped for 1 hour. The sample was analyzed under the same conditions again after 1 hour. The conventional C₁₈ column showed change of retention time, but COSMOSIL 5C₁₈-PAQ maintained stable retention time.

Applications


• Organic Acids




• Nucleobases and Nucleosides



• dNTPs



• Catecholamines




Applications

• Ascorbic Acid

COSMOSIL Application Data

Column: 5C₁₈-PAQ
 Column size: 4.6mm I.D.-250mm
 Mobile phase: 20mmol/l Phosphoric Acid
 Flow rate: 1.0 ml/min
 Temperature: 30°C
 Detection: UV245nm, 0.16AUFS
 Sample: L(+)-Ascorbic Acid [Vitamin C] (0.1μg)



NACALAI TESQUE, INC


AP-0372

• 2-Phenylethyl Glycosides

COSMOSIL Application Data

Column: 5C₁₈-PAQ
 Column size: 4.6mm I.D.-150mm
 Mobile phase: Acetonitrile/ Methanol/ H₂O = 8/4/8
 Flow rate: 1.0 ml/min
 Temperature: 30°C
 Detection: UV210nm

Sample: 1; 2-Phenylethyl-β-melibioside
 2; 2-Phenylethyl-β-gentiobioside
 3; 2-Phenylethyl-β-lactoside
 4; 2-Phenylethyl-β-cellobioside
 5; 2-Phenylethyl-β-maltoside



NACALAI TESQUE, INC

Data courtesy of Dr. K. Sakata, Dr. B. Shimizu, Institute for Chemical Research, Kyoto University

Ordering Information

• Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL 5C₁₈-PAQ Packed Column

Column Size I.D. x Length (mm)	Product Number
1.0 x 50	05792-61
1.0 x 150	05793-51
2.0 x 30	05878-51
2.0 x 50	05794-41
2.0 x 100	05470-71
2.0 x 150	34449-71
2.0 x 250	05795-31
3.0 x 100	05796-21
3.0 x 150	05797-11
3.0 x 250	05798-01
4.6 x 30	05879-41
4.6 x 50	34451-21

COSMOSIL 5C₁₈-PAQ Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	02484-91
10 x 20	34457-61
20 x 20	05803-11
20 x 50	05804-01
28 x 50	34455-81

• Preparative Columns (Particle Size: 15 μm)

COSMOSIL 15C₁₈-PAQ Packed Column


Column Size I.D. x Length (mm)	Product Number
28 x 250	05888-21
50 x 250	05890-71
50 x 500	05891-61

COSMOSIL 15C₁₈-PAQ Guard Column

Column Size I.D. x Length (mm)	Product Number
28 x 50	05887-31
50 x 50	05889-11


COSMOSIL C₁₈-EB

- 3 μ m C₁₈ column with reduced tailing and high resolution
 - Usable with simple mobile phases
- < Suitable Samples >
- For quality control of drugs
 - Compounds that induce peak tailing, such as basic compounds



Analysis of Basic Compounds


COSMOSIL 3C₁₈-EB uses a new end-capping method to reduce the number of residual silanol groups, which can cause peak tailing with basic compounds.




Analysis of Acidic Compounds and Metal Coordination Compounds

COSMOSIL 3C₁₈-EB utilizes a new end-capping method and high-purity silica gel to reduce tailing with metal coordination compounds.

• Acidic Compounds



• Metal Coordination Compounds



Ordering Information

- Analytical Columns (Particle Size: 3 μ m)

COSMOSIL 3C₁₈-EB Packed Column

Column Size I.D. x Length (mm)	Product Number
2.0 x 50	09794-21
2.0 x 75	09795-11
2.0 x 100	09796-01
2.0 x 150	09797-91
2.0 x 250	09798-81
3.0 x 50	09799-71
3.0 x 75	09800-21
3.0 x 100	09811-81

Column Size I.D. x Length (mm)	Product Number
3.0 x 150	09814-51
3.0 x 250	09827-91
4.6 x 50	09840-01
4.6 x 75*	09841-91
4.6 x 100*	09842-81
4.6 x 150*	09843-71
4.6 x 250	09844-61

COSMOSIL 3C₁₈-EB Guard Column






Column Size I.D. x Length (mm)	Product Number
2.0 x 10 Cartridge**	11892-74
4.6 x 10	09839-41
4.6 x 10 Cartridge**	11890-94

* Validated columns

** 2 cartridges included. Guard cartridge holder required; refer to page 71.


Reversed Phase Specialty Columns

Specifications

Packing Material	Cholester	PBr	π NAP	PYE	NPE
Silica Gel	High purity porous spherical silica				
Average Particle Size	2.5, 5 μ m	5 μ m	2.5, 5 μ m	5 μ m	
Average Pore Size	approx. 120 Å				
Specific Surface Area	approx. 300 m ² /g				
Bonded Phase Structure					
Bonded Phase	Cholesteryl group	Pentabromobenzyl group	Naphtylethyl group	Pyrenylethyl group	Nitrophenylethyl group
Bonding Type	Monomeric				
Main Interaction	Hydrophobic interaction Molecular shape selectivity	Hydrophobic interaction Dispersion force	Hydrophobic interaction π - π interaction	Hydrophobic interaction π - π interaction Dispersion force Molecular shape selectivity	Hydrophobic interaction π - π interaction Dipole-dipole interaction
End-Capping Treatment	Near-perfect treatment				
Carbon Content	approx. 20%	approx. 8%	approx. 11%	approx. 18%	approx. 9%
pH Range	2-7.5				
Features	•Usable under the same conditions as C ₁₈ •High molecular shape selectivity	•Separate hydrophilic compounds under reversed-phase conditions	•Stronger π - π interaction than phenyl columns	•Very strong π - π interaction	•Strong dipole-dipole interaction

Selectivity for positional isomers of dinitrobenzene

Different stationary phase exhibits different selectivity due to the use of forces that C₁₈ (hydrophobic interaction) does not have. By using these columns, you can achieve separation that cannot be done using only C₁₈.




COSMOSIL Cholester

- Cholesterol-bonded stationary phase
- Increased stereoselectivity and improved resolution for geometric isomers
- Usable under the same conditions as C₁₈


< Suitable Samples >

- Natural compounds, polyphenols, catechins, fat-soluble vitamins and flavones




Molecular Shape Selectivity

The stationary phase of Cholester has a very rigid structure and can distinguish different molecular shapes. Cholester retains planar triphenylene longer than non-planar *o*-terphenyl.




Hydrophobic Interaction

The below figure shows the comparison of hydrophobic interactions with competitor's C₁₈ columns. Cholester provides about the same hydrophobicity as alkyl group bonded types (C₁₈, C₃₀). It is not necessary to change the analytical conditions when replacing C₁₈ or C₃₀ columns with Cholester.



Applications


• Catechins




Applications (Continued)

2.5 μm particles yield better performance and shorter analysis time compared to 5 μm particles.

• Saikosaponins



• Water-Soluble Vitamins



Ordering Information

• Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL Cholester Packed Column

Column Size I.D. x Length (mm)	Product Number
1.0 x 150	05968-71
1.0 x 250	05969-61
2.0 x 30	08565-51
2.0 x 50	06352-91
2.0 x 100	06948-01
2.0 x 150	05971-11
2.0 x 250	05972-01
3.0 x 150	05973-91
3.0 x 250	05974-81

COSMOSIL Cholester Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 150*	05976-61
4.6 x 150 3 lots set*	07970-03
4.6 x 250*	05977-51
10 x 150	08011-91
10 x 250	05979-31
20 x 150	06088-71
20 x 250	05982-71
28 x 250	05985-41

* Validated Columns

COSMOSIL Cholester Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	05975-71
10 x 20	05978-41
20 x 20	05980-91
20 x 50	05981-81
28 x 50	05983-61

• Analytical (Particle Size: 2.5 μm)

COSMOSIL 2.5Cholester Packed Column


Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
2.0 x 50	09000-01	3.0 x 50	09049-91
2.0 x 75	09047-11	3.0 x 75	09050-51
2.0 x 100	09048-01	3.0 x 100	09051-41

COSMOSIL PBr

- Pentabromobenzyl-bonded stationary phase
- Separate hydrophilic compounds in reversed-phase conditions


< Suitable Samples >

- Hydrophilic compounds
- Nucleotides, peptides, catecholamines and oligosaccharides




Comparison with C₁₈

COSMOSIL PBr retains hydrophilic compounds stronger than C₁₈ columns under the same reversed-phase conditions.



Comparison with HILIC

HILIC is widely recognized as a method for separating hydrophilic compounds. However, because it differs from the commonly used reversed-phase mode, setting mobile phase conditions can be difficult. In addition, the use of acetonitrile in high concentration can cause problems with peak shape when using water as a sample solvent. COSMOSIL PBr can retain hydrophilic compounds under reversed-phase conditions, and maintains good peak shape even when injecting large amounts of water.



Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL PBr Packed Column

Column Size I.D. x Length (mm)	Product Number
2.0 x 100	13245-81
2.0 x 150	12392-81
2.0 x 250	13247-61
3.0 x 50	12592-61
3.0 x 100	13249-41
3.0 x 150	13250-01
3.0 x 250	13251-91
4.6 x 50	13252-81
4.6 x 150	12394-61
4.6 x 250	12395-51

COSMOSIL PBr Guard Column


Column Size I.D. x Length (mm)	Product Number
4.6 x 10 Catridge*	12444-14
10 x 20	12396-41
20 x 20	13256-41

* 2 cartridges included. Guard cartridge holder required; refer to page 71.


Column Size I.D. x Length (mm)	Product Number
10 x 50	13253-71
10 x 100	13254-61
10 x 150	13255-51
10 x 250	12397-31
20 x 50	13257-31
20 x 100	13258-21
20 x 150	13259-11
20 x 250	12398-21
28 x 100	13260-71
28 x 150	13261-61
28 x 250	13262-51

COSMOSIL π NAP

- Naphthalene-bonded stationary phase
 - Enhanced π - π interactions
- < Suitable Samples >
- Aromatic compounds and positional isomers



Comparison of π - π Interactions




COSMOSIL π NAP shows stronger π - π interactions than phenyl columns. Its two fused aromatic rings retain nitrobenzene stronger than phenyl columns.

Figure. Comparison of π - π interaction


Applications

• Adrenal Cortical Hormones



2.5 μ m particles yield better performance and shorter analysis time compared to 5 μ m particles.

• Berberine



Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 µm)

COSMOSIL πNAP Packed Column

Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
1.0 x 150	08076-61	3.0 x 250	08081-81
1.0 x 250	08077-51	4.6 x 150	08085-41
2.0 x 30	08566-41	4.6 x 250	08086-31
2.0 x 50	08567-31	10 x 150	08088-11
2.0 x 100	08299-51	10 x 250	08089-01
2.0 x 150	08078-41	20 x 150	08092-41
2.0 x 250	08079-31	20 x 250	08093-31
3.0 x 150	08080-91	28 x 250	08095-11

COSMOSIL πNAP Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	08082-71
10 x 20	08087-21
20 x 20	08090-61
20 x 50	08091-51
28 x 50	08094-21


- Analytical Columns (Particle Size: 2.5 µm)

COSMOSIL πNAP Packed Column


Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
2.0 x 50	06062-91	3.0 x 50	06054-01
2.0 x 75	06051-31	3.0 x 75	06055-91
2.0 x 100	06052-21	3.0 x 100	06057-71

COSMOSIL PYE

- Pyrenylethyl-bonded stationary phase
- Stronger π - π interactions
- < Suitable Samples >
- Aromatic compounds, positional isomers, dioxins, and PCBs



Comparison of π - π Interaction



COSMOSIL PYE provides much stronger π - π interactions than π NAP page 26.





Figure. Comparison of π - π interactions

Applications

- Sterols



Caution

1. Methanol is the recommended mobile phase for COSMOSIL PYE. Acetonitrile is not recommended because it has many π electrons and interferes with π - π interactions between the sample and the stationary phase.
2. The stationary phase of COSMOSIL PYE, pyrenylethyl group, has a large UV absorption. When the stationary phase detaches from silica gel and elutes, even a slight quantity can be detected and causes baseline noise. In such cases, wash the column with tetrahydrofuran. Detachment of a small amount of the stationary phase does not deteriorate a column's separation ability.
3. COSMOSIL PYE column is not suitable for gradient analysis.

Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μ m)

COSMOSIL 5PYE Packed Column

Column Size I.D. x Length (mm)	Product Number
1.0 x 150	02851-71
2.0 x 150	38042-61
2.0 x 250	34450-31

COSMOSIL 5PYE Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 150	37837-91
4.6 x 250	37989-11
10 x 250	37996-11
20 x 250	38044-41


Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37903-11
10 x 20	38041-71
20 x 20	05867-91
20 x 50	34475-21

COSMOSIL NPE

- Nitrophenylethyl-bonded stationary phase
- Separation with dipole-dipole and π - π interactions


< Suitable Samples >

- Isomers and nitro compounds



Selectivity for Dipole-Dipole Interactions

COSMOSIL NPE strongly retains 1,8-dinitronaphthalene because of the strong dipole formed by the two nitro groups positioned on the same side of naphthalene.



Caution

1. Methanol is recommended as a mobile phase for COSMOSIL NPE. Acetonitrile is not recommended because it has many π electrons and interferes with π - π interactions between the sample and the stationary phase.
2. The stationary phase of COSMOSIL NPE, nitrophenyl group, has a large UV absorption. When the stationary phase detaches from silica gel and elutes, even a slight quantity can be detected and causes baseline noise. In such a case, wash the column with tetrahydrofuran. Detachment of a small amount of the stationary phase does not deteriorate a column's separation ability.
3. COSMOSIL NPE column is not suitable for gradient analysis.

Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μ m)

COSMOSIL 5NPE Packed Column






Column Size I.D. x Length (mm)	Product Number
1.0 x 150	05897-01
2.0 x 150	34328-51
2.0 x 250	34379-91

COSMOSIL 5NPE Guard Column


Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
4.6 x 150	37902-21	4.6 x 10	37904-01		
4.6 x 250	37990-71	10 x 20	38045-31		
10 x 250	05469-11	20 x 20	05868-81		
20 x 250	38046-21	20 x 50	05869-71		

Other Reversed Phase Columns

Specifications

Packing Material	PFP	CN-MS	C ₂₂ -AR-II	C ₈ -MS	C ₄ -MS	TMS-MS	PE-MS
Silica Gel	High purity porous spherical silica						
Average Particle Size	5 μm						
Average Pore Size	approx. 120 Å						
Specific Surface Area	approx. 300 m ² /g						
Bonded Phase Structure							
Bonded Phase	Pentafluorophenyl group	Cyanopropyl group	Dococyl group	Octyl group	Butyl group	Trimethyl group	Phenylethyl group
Bonding Type	Monomeric		Polymeric	Monomeric			
Main Interaction	Hydrophobic interaction π-π interaction Dipole-dipole	Hydrophobic interaction π-π interaction	Hydrophobic interaction				Hydrophobic interaction π-π interaction
End-Capping Treatment	Near-perfect treatment						
Carbon Content	approx. 10%	approx. 7%	approx. 19%	approx. 10%	approx. 7%	approx. 5%	approx. 10%
pH Range	2-7.5						

Difference in Separation Characteristics



COSMOSIL PFP



- Pentafluorophenyl-bonded stationary phase
- Alternative selectivity to C₁₈ columns

< Suitable Samples >

- Vitamin E, structural isomers and fluorides


Alternative Selectivity to C₁₈ Columns

COSMOSIL PFP provides different selectivity from C₁₈ Columns. Furthermore, it offers improved separation compared to other PFP columns.




Applications

• Vitamin E



• Cresol Isomers



Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL 5PFP Packed Column

Column Size I.D. x Length (mm)	Product Number	Column Size I.D. x Length (mm)	Product Number
2.0 x 50	13263-41	10 x 50	13272-21
2.0 x 100	13264-31	10 x 100	13273-11
2.0 x 150	12381-21	10 x 150	13274-01
2.0 x 250	13265-21	10 x 250	12386-71
3.0 x 50	13266-11	20 x 50	13276-81
3.0 x 100	13267-01	20 x 100	13277-71
3.0 x 150	13268-91	20 x 150	13278-61
3.0 x 250	13269-81	20 x 250	12387-61
4.6 x 50	13270-41	28 x 100	13280-11
4.6 x 100	13271-31	28 x 150	13281-01
4.6 x 150	12383-01	28 x 250	13282-91
4.6 x 250	12384-91		

COSMOSIL 5PFP Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10 Cartridge*	12443-24
10 x 20	12385-81
20 x 20	13275-91
28 x 50	13279-51


* 2 cartridges included. Guard cartridge holder required; refer to page 71.

COSMOSIL CN-MS


- Cyanopropyl-bonded stationary phase
- Enables separation of different hydrophobic samples without using gradient
- < Suitable Samples >
- Mixtures of natural compounds

Rapid Analysis

Gradient elution is commonly used for the samples containing both polar and non-polar compounds. However, gradient elution may cause reproducibility problems depending on the gradient mixer and pump, and needs equilibration time for each analysis. COSMOSIL 5CN-MS offers rapid analysis and great reproducibility using isocratic elution mode.



Application of Different Hydrophobic Samples




Applications

- Acetoaminophen


COSMOSIL Application Data

Column:
Column size: 4.6mm I.D.-150mm
Mobile phase: Methanol/ 50mmol/l KH_2PO_4
(pH4.7 with NaOH) = 20/80
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV225nm

Sample: 1; *p*-Aminophenol (0.2μg)
2; *p*-Acetamidophenol (0.2μg)
3; 4'-Acetoxyacetanilide (0.2μg)



5CN-MS



NACALAI TESQUE, INC

AP-1075

Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL 5CN-MS Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	38233-61
4.6 x 100	38234-51
4.6 x 150	38235-41
4.6 x 250	38236-31

COSMOSIL 5CN-MS Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	38231-81
10 x 20	38232-71

Column Size I.D. x Length (mm)	Product Number
6.0 x 150	38237-21
6.0 x 250	38238-11
10 x 250	38239-01
20 x 250	38240-61

COSMOSIL C₂₂-AR-II, C₈-MS, C₄-MS, TMS-MS, PE-MS

Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 µm)

COSMOSIL 5C₂₂-AR-II Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	05848-41
4.6 x 100	05849-31
4.6 x 150	04598-51
4.6 x 250	04599-41

COSMOSIL 5C₂₂-AR-II Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	04881-21
10 x 20	05554-81

COSMOSIL 5C₈-MS Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	38153-11
4.6 x 100	38154-01
4.6 x 150	38155-91
4.6 x 250	38156-81

COSMOSIL 5C₈-MS Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	38151-31
10 x 20	38152-21

COSMOSIL 5C₄-MS Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	38163-81
4.6 x 100	38164-71
4.6 x 150	38165-61
4.6 x 250	38166-51

COSMOSIL 5C₄-MS Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	38161-01
10 x 20	38162-91

COSMOSIL 5TMS-MS Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	38173-51
4.6 x 100	38174-41
4.6 x 150	38175-31
4.6 x 250	38176-21

COSMOSIL 5TMS-MS Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	38171-71
10 x 20	38172-61

COSMOSIL 5PE-MS Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	38183-21
4.6 x 100	38184-11
4.6 x 150	38185-01
4.6 x 250	38186-91

COSMOSIL 5PE-MS Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	38181-41
10 x 20	38182-31

(2) Normal Phase Columns

COSMOSIL SL-II


- High purity silica gel (>99.99%) with special treatment
- Suitable for preparative separation

Specifications

Packing Material	SL-II
Silica Gel	High purity porous spherical silica
Average Particle Size	3, 5, 15 μm
Average Pore Size	approx. 120 Å
Specific Surface Area	approx. 300 m ² /g
Features	• High purity silica gel (>99.99%) with special treatment • Suitable for preparative separation (higher resolution than medium-pressure or open chromatography)

Comparison with Old Type

COSMOSIL SL-II with high purity silica gel offers better peak shape for phenols with a simple mobile phase of ethanol or hexane. No acetic acid additives were required, unlike for the old type silica.



Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL 5SL-II Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	37999-81
4.6 x 100	38000-01
4.6 x 150	38001-91
4.6 x 250	38002-81

Column Size I.D. x Length (mm)	Product Number
6.0 x 150	38003-71
6.0 x 250	38004-61
10 x 250	38005-51
20 x 250	38006-41
28 x 250	34358-61

COSMOSIL 5SL-II Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37997-01
10 x 20	37998-91
20 x 20	05874-91
20 x 50	05875-81
28 x 50	34359-51

- Preparative Columns (Particle Size : 15 μm)

COSMOSIL 15SL-II Packed Column

Column Size I.D. x Length (mm)	Product Number
28 x 250	05893-41
50 x 250	05895-21
50 x 500	05896-11

COSMOSIL 15SL-II Guard Column

Column Size I.D. x Length (mm)	Product Number
28 x 50	05892-51
50 x 50	05894-31

- Fast LC column (Particle Size: 3 μm)

COSMOSIL 3SL-II Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	38059-61
4.6 x 50	38060-21
4.6 x 100	38061-11

(3) Hydrophilic Interaction Columns

COSMOSIL HILIC

- Triazole bonded stationary phase
- Enhanced hydrophilic interaction
- Unique anion-exchange mechanism

< Suitable Samples >


- Hydrophilic compounds that would not be retained in reversed phase chromatography
- Melamine and water-soluble vitamins

Specifications

Packing Material	HILIC
Silica Gel	High purity porous spherical silica
Average Particle Size	2.5, 5 μm
Average Pore Size	approx. 120 Å
Specific Surface Area	approx. 300 m ² /g
Bonded Phase	Triazole
Interaction	Hydrophilic interaction, anion exchange
Target Substance	Hydrophilic compounds, acidic compounds
Features	Suitable for compounds not retained by C ₁₈

Comparison with C₁₈


The hydrophilic interaction chromatography is a variation of normal phase chromatography where a polar stationary phase is used with a mobile phase which contains a high concentration of water-miscible organic solvent and a low concentration of aqueous eluent. The main retention mechanism is the partitioning of the polar analytes between the polar stationary and the non-polar mobile phase. As it is also called “aqueous normal phase”, the elution order is similar to that of normal phase and the sample elution is in the order of increasing hydrophilicity. Without using ion-pair reagent COSMOSIL HILIC retains highly polar analytes that would not be retained in reversed phase chromatography. It also shows a weak anion-exchange mechanism with the positively charged stationary phase, thus acidic compounds are strongly retained.




Applications

2.5 μ m particles yield better performance and shorter analysis time compared to 5 μ m particles.


• Nucleotides




• Phosphorylated Peptide



• Umami Components



• Melamine



Ordering Information

• Analytical / Preparative Columns (Particle Size: 5 μ m)

COSMOSIL HILIC Packed Column

Column Size I.D. x Length (mm)	Product Number
1.0 x 150	07869-11
1.0 x 250	07870-71
2.0 x 30	08568-21
2.0 x 50	07052-91
2.0 x 100	08569-11
2.0 x 150	07054-71
2.0 x 250	07489-91
3.0 x 150	07871-61
3.0 x 250	07872-51

Column Size I.D. x Length (mm)	Product Number
4.6 x 150*	07056-51
4.6 x 150 3 lots set	09385-23
4.6 x 250*	07057-41
10 x 250	07059-21
20 x 250	07060-81
28 x 250	07875-21

* Validated Columns

COSMOSIL HILIC Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	07055-61
10 x 20	07058-31
20 x 20	07854-91
20 x 50	07873-41
28 x 50	07874-31

• Analytical Columns (Particle Size: 2.5 μ m)

COSMOSIL HILIC Packed Column

Column Size I.D. x Length (mm)	Product Number
2.0 x 50	11766-21
2.0 x 75	11768-01
2.0 x 100	11769-91
2.0 x 150	11770-51


Column Size I.D. x Length (mm)	Product Number
3.0 x 50	11771-41
3.0 x 75	11772-31
3.0 x 100	11773-21
3.0 x 150	11773-21

(4) Mono- and Oligosaccharide Analysis Columns

Introduction


Saccharides are not retained on standard C₁₈ columns because of their low hydrophobicity. COSMOSIL Sugar-D and NH₂-MS are specifically designed for separation of saccharides. COSMOSIL C₁₈-PAQ is recommended for hydrophobic glycosides or saccharide derivatives.

Specifications

Packing Material	Sugar-D	NH ₂ -MS
Silica Gel	High purity porous spherical silica	
Average Particle Size	5 μm	
Average Pore Size	—	approx. 120 Å
Specific Surface Area	—	approx. 300 m ² /g
Bonded Phase Structure	—	
Bonded Phase	Secondary/tertiary amine	Aminopropyl group
Bonding Type	—	Polymeric
Target Substances	Monosaccharides, oligosaccharides	
End-Capping Treatment	—	Near-perfect treatment
Carbon Content	—	approx. 4%
Features	•First choice for saccharide analysis •High durability •Good quantitative analysis	•Different selectivity from Sugar-D

Comparison of Retention

The conventional aminopropyl column is slightly more retentive than Sugar-D. The retention time can be adjusted by increasing the concentration of acetonitrile in the mobile phase by 5%-10% as shown below.




COSMOSIL Sugar-D

- Novel stationary phase for saccharides
- Superior durability compared to conventional amino columns
- Minimized undesirable adsorption

Comparison of Durability

The decrease of retention time was compared between COSMOSIL Sugar-D and conventional aminopropyl bonded stationary phase with a severe 100% water eluent between tests. The capacity factor of Sugar-D did not decrease.




Decomposition Condition

Solution	Water
Flow Rate	1.0 ml/min
Temperature	Room Temperature
Column	4.6 mm I.D. x 250 mm
Mobile Phase	Acetonitrile : Water = 70 : 30
Flow Rate	1.0 ml/min
Temperature	30°C
Detection	RI
Sample	Maltose


Comparison of Adsorption

Certain types of saccharides, such as arabinose or galactose, are partially or temporarily adsorbed on conventional aminopropyl stationary phases, causing tailing or no elution at all. COSMOSIL Sugar-D provides superior separation and high recovery for these saccharides.




Applications

• Mono- and Oligosaccharides



• Polyols



Applications


• Oligomaltoses

COSMOSIL Application Data

Column: Sugar-D
Column size: 4.6mm I.D.-250mm
Mobile phase: Acetonitrile / H₂O = 65/35
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: RI

Sample:

1; D-(+)-Glucose (10µg)
2; Maltose (10µg)
3; Maltotriose (10µg)
4; Maltotetraose (10µg)
5; Maltopentaose (10µg)
6; Maltohexaose (10µg)
7; Maltotriose (10µg)



NACALAI TESQUE, INC

AP-0331


• Oligofructoses

COSMOSIL Application Data

Column: Sugar-D
Column size: 4.6mm I.D.-250mm
Mobile phase: Acetonitrile / H₂O = 70/30
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: RI

Sample:

Fructooligosaccharides (50µg)
1; I-Kestose
2; Nystose
3; 1-Fructofuranosyl-D-nystose



NACALAI TESQUE, INC

AP-0320

Ordering Information

• Analytical / Preparative Columns (Particle Size : 5 µm)

COSMOSIL Sugar-D Packed Column

Column Size I.D. x Length (mm)	Product Number
2.0 x 250	05689-31
3.0 x 150	05690-91
3.0 x 250	05691-81

COSMOSIL Sugar-D Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	05394-81
10 x 20	05696-31
20 x 50	05694-51

Column Size I.D. x Length (mm)	Product Number
4.6 x 150	05395-71
4.6 x 250	05397-51
10 x 250	05692-71
20 x 250	05693-61

COSMOSIL NH₂-MS

- Aminopropyl-bonded stationary phase
- Different selectivity from Sugar-D

Comparison of Adsorption


NH₂-MS offers better separation than Sugar-D for some samples.

Comparison of Separation Property

Column:
Column size: 4.6mm I.D.-250mm
Mobile phase: (NH₂-MS) Acetonitrile / H₂O = 75/25
(Sugar-D) Acetonitrile / H₂O = 80/20
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: RI

Sample:

1; Sucrose (10 µg)
2; Maltose (10 µg)
3; Lactose (10 µg)
4; D-(+)-Trehalose (10 µg)



NACALAI TESQUE, INC

Ordering Information

• Analytical / Preparative Columns (Particle Size: 5 µm)

COSMOSIL 5NH₂-MS Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 150	38245-11
4.6 x 250	38246-01


Column Size I.D. x Length (mm)	Product Number
10 x 250	38249-71
20 x 250	38250-31

COSMOSIL 5NH₂-MS Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	38241-51
10 x 20	38242-41
20 x 50	06093-91

(5) Protein Separation Columns

Protein separation with HPLC



Reversed Phase Columns

COSMOSIL Protein-R

- Excellent separation
- High recovery rate
- Outstanding stability at low pH


Specifications

Packing Material	Protein-R
Silica Gel	High purity porous spherical silica
Average Particle Size	5 µm
Average Pore Size	approx. 300 Å
Specific Surface Area	approx. 150 m ² /g
Bonded Phase	Octadecyl group
Bonding Type	Polymeric
Main Interaction	Hydrophobic interaction
End-Capping Treatment	Near-perfect treatment
pH Range	1.5-7.5*
Features	• High recovery rate • Acid-resistant

*Optimal pH range of silica-based columns is between 2 and 7.5. Extreme pH may significantly decrease column lifetime.

Comparison of Separation

Protein-R shows sharper peaks for proteins than conventional C₁₈ wide-pore columns.



Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL Protein-R Packed Column

Column Size I.D. x Length (mm)	Product Number
2.0 x 150	06514-71
4.6 x 50	06525-31
4.6 x 150	06526-21
4.6 x 250	06527-11

COSMOSIL Protein-R Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	06518-31
10 x 20	06528-01
20 x 20	08692-81

COSMOSIL C₁₈-AR-300, C₈-AR-300, C₄-AR-300, Ph-AR-300

- Wide-pore reversed-phase column
- 4 types of phases (octadecyl, octyl, butyl and phenyl)


Specifications

Packing Material	5C ₁₈ -AR-300	5C ₈ -AR-300	5C ₄ -AR-300	5Ph-AR-300
Silica Gel	High purity porous spherical silica			
Average Particle Size	5 μm			
Average Pore Size	300 Å			
Specific Surface Area	150 m ² /g			
Bonded Phase Structure				
Bonded Phase	Octadecyl group	Octyl group	Butyl group	Phenyl group
Bonding Type	Polymeric			
Main Interaction	Hydrophobic interaction			Hydrophobic interaction π-π interaction
End-Capping Treatment	Near-perfect treatment			
pH Range	1.5-7.5*			
Carbon Content	approx. 12%	approx. 7%	approx. 6%	approx. 7%

*Optimal pH range of silica-based columns is between 2 and 7.5. Extreme pH may significantly decrease column lifetime.

Comparison of Separation

COSMOSIL AR-300 packed column series offers 3 types of alkyl phases and a phenyl phase.



Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL 5C₁₈-AR-300 Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	37911-01
4.6 x 150	37913-81
4.6 x 250	37914-71

COSMOSIL 5C₁₈-AR-300 Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37910-11
10 x 20	37965-11

COSMOSIL 5C₈-AR-300 Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	37951-81
4.6 x 150	37953-61
4.6 x 250	37954-51

Column Size I.D. x Length (mm)	Product Number
10 x 150	34345-21
10 x 250	34247-11
20 x 150	05861-51
20 x 250	34364-71

COSMOSIL 5C₈-AR-300 Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37950-91
10 x 20	34464-61

COSMOSIL 5C₄-AR-300 Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	37956-31
4.6 x 150	37958-11
4.6 x 250	37959-01

Column Size I.D. x Length (mm)	Product Number
10 x 150	34249-91
10 x 250	38047-11
20 x 150	34477-01
20 x 250	38048-01

COSMOSIL 5C₄-AR-300 Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37955-41
10 x 20	05862-41

COSMOSIL 5Ph-AR-300 Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	37961-51
4.6 x 150	37963-31
4.6 x 250	37964-21

Column Size I.D. x Length (mm)	Product Number
10 x 150	05865-11
10 x 250	34267-51
20 x 150	05866-01
20 x 250	34468-21

COSMOSIL 5Ph-AR-300 Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37960-61
10 x 20	34268-41

COSMOSIL Diol-120-II, Diol-300-II, Diol-1000-II

- Ideal for the size-based separation of proteins and water-soluble polymers
- Reduce undesirable adsorption

Specifications


Packing Material	5Diol-120-II	5Diol-300-II	5Diol-1000-II
Silica Gel	High purity porous spherical silica ⁽¹⁾		
Average Particle Size	5 µm		
Average Pore Size	approx. 120 Å	approx. 300 Å	approx. 1000 Å ⁽²⁾
Bonded Phase	Diol group		
Target Substances	Proteins, water soluble polymers		
Flow Rate	0.5-1.0 (ml/min)		
Selection of Pore Size (protein)	MW 5,000-100,000	MW 10,000-700,000	-
Selection of Pore Size (water-soluble polymers)	MW 1,000-20,000	MW 5,000-100,000	MW 50,000-500,000

(1) With the silica-based gel, organic solvents, including methanol and acetonitrile, can be used.

(2) If you require pore sizes greater than 1000 Å, please contact us.

Calibration Curve


• Calibration Curve of Proteins



Column COSMOSIL 5Diol-II (7.5 mm I.D. x 600 mm)
 Mobile Phase 20mmol/l Phosphate Buffer (pH7.0)+100mmol/l Na₂SO₄
 Flow Rate 1.0ml/min
 Temperature 30°C

Sample	M.W.	Sample	M.W.
Thyroglobulin	660,000	Peroxidase	40,000
Catalase	250,000	Carbonic Anhydrase	30,000
Glucose Oxidase	160,000	α-Chymotrypsinogen A	25,700
Uricase	128,000	α-Chymotrypsin	25,200
Choline Oxidase	95,000	Trypsinogen	24,000
Transferrin	85,000	Trypsin (bovine)	23,300
Conalbumin	77,500	Myoglobin	17,000
Malate Dehydrogenase	70,000	Lysozyme	14,300
α-Glucosidase	68,500	Ribonuclease A	13,700
Albumin (BSA)	66,000	Cytochrome C	12,400
α-Amylase	52,500	Aprotinin	6,500
Fetuin	48,000	Gly-Gly	132
Albumin (Ovalbumin)	45,000		

• Linear pullulan calibration curve




Column COSMOSIL 5Diol-II (7.5 mm I.D. x 300 mm)
 Mobile Phase Water
 Flow Rate 1.0ml/min
 Temperature 30°C
 Detection RI
 Sample Linear pullulan

Sample	M.W.
1; P-800	853,000
2; P-400	380,000
3; P-200	186,000
4; P-100	100,000
5; P-50	48,000
6; P-20	23,700
7; P-10	12,200
8; P-5	5,800
9; Maltotriose	504
10; Maltose	342
11; Glucose	180


Molecular Weight Range

Globular molecules have smaller apparent size compared to linear molecules of the same weight. Therefore, globular molecules can be separated with smaller pores compared to linear molecules.



Application

• DNA



Ordering Information

• Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL 5Diol-120-II Packed Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 300	38050-51
7.5 x 600	38051-41

COSMOSIL 5Diol-120-II Guard Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 50	38049-91

COSMOSIL 5Diol-300-II Packed Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 300	38053-21
7.5 x 600	38054-11

COSMOSIL 5Diol-300-II Guard Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 50	38052-31

COSMOSIL 5Diol-1000-II Packed Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 300	13338-71

COSMOSIL 5Diol-1000-II Guard Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 50	13337-81

COSMOGEL IEX Series

- Available in 3 different ion-exchange modes (Anion-exchange type, Cation-exchange type, Amphoteric ion-exchange type)
- Available for 3 different application areas (for Purification, for Ultra-fast analysis, for Precise analysis)
- For separation of biopolymers such as proteins or nucleic acids

Specifications

Packing Material	Type Q	Type Q-N	Type S	Type S-N	Type M	Type M-N
Gel	Hydrophilic polymer					
Average Particle Size	5 μ m					
Average Pore Size	1000 Å	Non-porous	1000 Å	Non-porous	1000 Å	Non-porous
Functional Group	$-\text{CH}_3\text{N}^+(\text{CH}_3)_3$		$-(\text{CH}_2)_3\text{SO}_3^-$		$-\text{CH}_3\text{N}^+(\text{CH}_3)_3 + -(\text{CH}_2)_3\text{SO}_3^-$	
Protein Binding Capacity	110-150 mg	12-20 mg	70-100 mg	10-18 mg	55-75 mg(BSA)/ml 35-50 mg(IgG)/ml	6-10 mg(BSA)/ml 5-9 mg(IgG)/ml
Column Size I.D. x Length (mm)	4.6 x 50	4.6 x 30 4.6 x 100	4.6 x 50	4.6 x 30 4.6 x 100	4.6 x 50	4.6 x 100
Column Material	PEEK					
Connection	Waters Type					


Type of Packing Material

COSMOGEL IEX Series are available in amphoteric ion-exchange type in which two kinds of packing materials are mixed, as well as in widely used anion-exchange type and cation-exchange type.

Type of Packing Material	Target Sample	Average Pore Size	
		Porous (1000 Å)	Non-porous
Anion-Exchange Type	Acidic proteins / DNA	Type Q	Type Q-N
Cation-Exchange Type	Basic proteins	Type S	Type S-N
Amphoteric Ion-Exchange Type	All proteins	Type M	Type M-N

Comprehensive isolation of proteins by amphoteric ion-exchange type (Type M)

The amphoteric ion-exchange type enables the simultaneous separation of both acidic and basic proteins in one application.



Type of Column

COSMOGEL IEX columns are available for 3 types of applications:



Application	Pore Size	Column Size I.D. x Length (mm)	Column		
For Purification	Porous (1000 Å)	4.6 × 50	Type Q	Type S	Type M
For Precise Analysis	Non-porous	4.6 × 100	Type Q-N	Type S-N	Type M-N
For Ultra-Fast Analysis	Non-porous	4.6 × 30	Type Q-N	Type S-N	—

For Purification: Type Q, Type S, Type M

Porous packing materials have higher binding capacity for proteins than the non-porous type, which means that peak shape does not spread even with injection of a large volume of sample. Therefore they are highly suitable for purification of large samples.

COSMOSIL Application Data

Column: COSMOGEL IEX Type S
 Column size: 4.6mm I.D. x 50mm
 Mobile phase: A: 20mmol/l Phosphate buffer(pH6.8)
 B: 20mmol/l Phosphate buffer(pH6.8) + 0.5mol/l NaCl
 B conc. 0→100% 30min Linear gradient
 Flow rate: 0.5 ml/min
 Temperature: 30°C
 Detection: UV280nm
 Sample: 1; Ribonuclease A
 2; Cytochrome C
 3; Lysozyme



NACALAI TESQUE, INC


AP-1185

For Precise Analysis: Type Q-N, Type S-N, Type M-N

Non-porous packing materials reduce spreading of samples in packing materials, resulting in high resolution separation for precise analysis, such as quality control of antibody preparations. The longer column length also contributes to the sharper peaks.

COSMOSIL Application Data

Column: COSMOGEL IEX
 Column size: 4.6mm I.D. x 50mm (Type S) / 4.6mm I.D. x 100mm (Type S-N)
 Mobile phase: A: 20mmol/l Phosphate buffer(pH6.8)
 B: 20mmol/l Phosphate buffer(pH6.8) + 0.5mol/l NaCl
 B conc. 0→100% 30min Linear gradient
 Flow rate: 0.5 ml/min
 Temperature: 30°C
 Detection: UV220nm
 Sample: 1; Ribonuclease A (1.65 µg)
 2; Cytochrome C (1.65 µg)
 3; Lysozyme (1.65 µg)



NACALAI TESQUE, INC


AP-1181

For Ultra-fast Analysis: Type Q-N, Type S-N

Non-porous packing materials are not much affected by high flow rate and thus the materials are suitable for fast analysis. The shorter column length contributes to the fast analysis.

COSMOSIL Application Data

Column: COSMOGEL IEX
 Column size: 4.6mm I.D. x 50mm (Type S) / 4.6mm I.D. x 30mm (Type S-N)
 Mobile phase: A: 20mmol/l Phosphate buffer(pH6.8)
 B: 20mmol/l Phosphate buffer(pH6.8) + 0.5mol/l NaCl
 B conc. 0→100% Linear gradient
 Type S: 30min
 Type S-N: 10min
 Flow rate: Type S: 0.5 ml/min
 Type S-N: 1.0 ml/min
 Temperature: 30°C
 Detection: UV220nm
 Sample: 1; Ribonuclease A (1.65 µg)
 2; Cytochrome C (1.65 µg)
 3; Lysozyme (1.65 µg)



NACALAI TESQUE, INC

AP-1183

Ordering Information

Ion Exchange Mode	Product Name	Application	Column Size I.D. x Length (mm)	Product Number
Anion-exchange Type	COSMOGEL IEX Type Q	For Purification	4.6 x 50	06266-31
	COSMOGEL IEX Type Q-N	For Ultra-fast Analysis	4.6 x 30	06264-51
	COSMOGEL IEX Type Q-N	For Precise Analysis	4.6 x 100	06258-41
Cation-exchange Type	COSMOGEL IEX Type S	For Purification	4.6 x 50	06252-01
	COSMOGEL IEX Type S-N	For Ultra-fast Analysis	4.6 x 30	06251-11
	COSMOGEL IEX Type S-N	For Precise Analysis	4.6 x 100	06250-21
Amphoteric Ion-exchange Type	COSMOGEL IEX Type M	For Purification	4.6 x 50	06248-71
	COSMOGEL IEX Type M-Nz	For Precise Analysis	4.6 x 100	06244-11

COSMOSIL HIC

- Separate based on differences in hydrophobicity
- Little loss in enzyme activity and the tertiary structure of proteins


Specifications

Packing Material	HIC
Silica Gel	High purity porous spherical silica
Average Particle Size	5 μm
Average Pore Size	approx. 300 $\text{\AA}$
Specific Surface Area	approx. 150 m^2/g
Main Interaction	Hydrophobic interaction


Applications

A buffer with high salt concentration, usually 1-2 mol/l of $(\text{NH}_4)_2\text{SO}_4$, is used as an initial mobile phase for adsorption of samples to a weakly hydrophobic stationary phase. The elution is done with a decreasing salt gradient. The application in the lower left shows that myoglobin elutes earlier than BSA under the buffer with high salt concentration, suggesting that myoglobin is less hydrophobic than BSA.

• Separation of Protein Standards



• Separation of β -Glucosidase



Ordering Information

- Analytical Columns (Particle Size: 5 μm)

COSMOSIL 5HIC Packed Column







Column Size I.D. x Length (mm)	Product Number
4.6 x 50	04263-21

(6) Columns for Fullerene Separation

Introduction


Separation of fullerenes, especially preparative scale separation, on conventional HPLC columns is always problematic due to the low solubility and low recovery rate of fullerenes. COSMOSIL offers a variety of columns designed for preparative scale separation of fullerenes, including higher fullerenes, metallofullerenes and fullerene derivatives.

Specifications

Packing Material	Buckyprep	Buckyprep-D	Buckyprep-M	PBB	PYE	NPE
Silica Gel	High purity porous spherical silica					
Average Particle Size	5 μm					
Average Pore Size	approx. 120 $\text{\AA}$					
Specific Surface Area	approx. 300 m^2/g					
Bonded Phase Structure						
Bonded Phase	Pyrenylpropyl group	Nitro-carbazoyl group	Phenothiazinyl group	Pentabromobenzyl group	Pyrenylethyl group	Nitrophenylethyl group
Bonding Type	Monomeric					
End-Capping Treatment	Near-perfect treatment		None	Near-perfect treatment		
Carbon Content	approx. 17%	-	approx. 13%	approx. 8%	approx. 18%	approx. 9%
Features	• Standard column for fullerene separation.	• For separation of derivatized fullerenes	• Designed to separate metallofullerenes	• Designed for preparative separation of C_{60} , C_{70}	• Separation of fullerene and structural isomers	• Separation of fullerene derivatives

Comparison of Retention

The figure below shows the retention time of C_{60} and C_{70} in toluene. COSMOSIL fullerene separation columns (Buckyprep, Buckyprep-D, Buckyprep-M, PBB and PYE) exhibit high fullerene retention with toluene, so they can easily separate C_{60} and C_{70} .



Suggested Solvents for Fullerene Separation


Solvent	Solubility of C_{60} (mg/ml)	Features	Solvent	Solubility of C_{60} (mg/ml)	Features
Toluene	3.2	The most commonly used solvent.	Chlorobenzene	7.0	Stronger eluent than toluene. Recommended for higher fullerenes.
<i>n</i> -Hexane	0.046	Weaker eluent than toluene	<i>o</i> -Dichlorobenzene	27.0	Stronger eluent than chlorobenzene.
<i>n</i> -Heptane	--		1,2,4-Trichlorobenzene	21.3	Strongest eluent. Recommended as a washing solvent.
Methanol	0.001				
2-Propanol	--				
Acetonitrile	0.018	Weaker eluent than toluene. Recommended as a washing solvent for Buckyprep-D.			

COSMOSIL Buckyprep

- Standard column for fullerene separation
- Excellent separation for higher and derivatized fullerenes


Difference in Preparative Separation

Buckyprep can be used with toluene, the most commonly-used solvent in fullerene separation. Because tailing does not occur, you can inject about 35 times more sample than with a C₁₈ column.




Applications

• Higher Fullerenes



• Oxidized Fullerenes



Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL Buckyprep Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 250	37977-61
10 x 250	37981-91
20 x 250	37982-81
28 x 250	34346-11

COSMOSIL Buckyprep Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37983-71
10 x 20	37984-61
20 x 50	34374-41
28 x 50	05871-21

COSMOSIL Buckyprep-D


- For preparative separation of derivatized fullerenes
- For separation of derivatized fullerenes such as C₆₀-indene (used for organic thin-film solar cell)

Applications

Buckyprep-D offers improved separation for C₆₀-indene, a derivatized fullerene that has received much attention as an n-type semiconductor material for organic thin-film solar cells.

COSMOSIL Application Data

Column:
Column size: 4.6mm I.D.-250mm
Mobile phase: Toluene
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV 325nm
Sample: C₆₀ [Indene]₂ (1.0mg/ml)
Inj.Vol.: 1.0μl




Data courtesy of Yusuke Tajima, Dr. Sci.
Organic Optoelectronics Laboratory, RIKEN (Institute of Physics and Chemistry)

NACALAI TESQUE, INC

Buckyprep-D retains derivatized fullerenes longer than C₆₀. Therefore it is more suitable for preparative separation of derivatized fullerenes than our conventional Buckyprep column.

COSMOSIL Application Data

Column:
Column size: 4.6mm I.D.-250mm
Mobile phase: Toluene
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV 325nm
Sample: 1; C₆₀ (0.25mg/ml)
2; [6,6]-Phenyl-C₆₁ Butyric Acid Methyl Ester [PCBM] (0.25mg/ml)
Inj.Vol. 1.0μl



NACALAI TESQUE, INC

Note

The baseline of Bukyprep-D is less stable relative to other fullerene columns. To stabilize baseline, let acetonitrile run through for 10 minutes before analysis.

Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL Buckyprep-D Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 50	09646-61
4.6 x 250	09647-51
10 x 250	09650-91
20 x 250	09651-81

COSMOSIL Buckyprep-D Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	09611-01
10 x 20	09613-81
20 x 50	09614-71

COSMOSIL Buckyprep-M

- Different selectivity from Buckyprep
- Excellent separation for metallofullerenes

Applications


Metallofullerenes

COSMOSIL Buckyprep-M is a phenothiazinyl-bonded silica-based column specifically designed for metallofullerene separation. Metallofullerenes are retained more strongly than other fullerenes on this column. COSMOSIL Buckyprep-M is also effective for the separation of higher fullerenes and fullerene derivatives.


COSMOSIL Application Data

Column: 4.6mm I.D. x 250mm
Mobile phase: Toluene
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV312nm

Sample: Sc₂@C₇₆(I)
Sc₂@C₇₈
Sc₂@C₈₀(I)
C₈₆



Sample courtesy of Prof. H.Shinohara, Department of chemistry, Nagoya University




NACALAI TESQUE, INC


COSMOSIL Application Data

Column: 4.6mm I.D. x 250mm
Mobile phase: Toluene
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV312nm

Sample: Gd@C₈₂(I)



Sample courtesy of Prof. H.Shinohara, Department of chemistry, Nagoya University



NACALAI TESQUE, INC

Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 µm)

COSMOSIL Buckyprep-M Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 250	04138-71
10 x 250	04141-11
20 x 250	04142-01
28 x 250	05873-01

COSMOSIL Buckyprep-M Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	04139-61
10 x 20	04140-21
20 x 50	34474-31
28 x 50	05872-11

COSMOSIL PBB

- Can be used with *o*-dichlorobenzene or carbon disulfide
- Suitable for preparative scale separation


Applications

The loading capacity of COSMOSIL PBB for C₆₀ and C₇₀ can be three times greater than COSMOSIL Buckyrep.

Preparative Separation of Fullerenes

Column:
Column size: 20mm I.D.-250mm
Mobile phase: Toluene
Flow rate: 18 ml/min
Temperature: Room temperature
Detection: UV285nm

Sample: Crude fullerenes (3.5mg/ml)



Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 µm)

COSMOSIL 5PBB Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 250	37980-01
10 x 250	37985-51
20 x 250	37986-41

COSMOSIL 5PBB Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37987-31
10 x 20	37988-21
20 x 50	34375-31

COSMOSIL NPE

- Different selectivity from Buckyrep or PBB
- Excellent separation for derivatized fullerenes

Applications


- PCBM, PCBB

COSMOSIL NPE retains derivatized C₆₀ stronger than C₆₀.

COSMOSIL Application Data

Column:
Column size: 4.6mm I.D.-250mm
Mobile phase: (Buckyrep, PBB) Toluene
(NPE) Toluene/Hexane = 25/75
Flow rate: 1.0 ml/min
Temperature: 30°C
Detection: UV325nm

Sample:
1; [6,6]-Phenyl-C₆₁ Butyric Acid Methyl Ester [PCBM] (1.5µg)
2; [6,6]-Phenyl-C₆₁ Butyric Acid Butyl Ester [PCBB] (1.5µg)
3; C₆₀ (1.5µg)



Hexane added to mobile phase due to NPE's weak retention.

Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL 5NPE Packed Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 150	37902-21
4.6 x 250	37990-71
10 x 250	05469-11
20 x 250	38046-21

COSMOSIL 5NPE Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37904-01
10 x 20	38045-31
20 x 50	05869-71

COSMOSIL PYE

Ordering Information

- Analytical / Preparative Columns (Particle Size: 5 μm)

COSMOSIL 5PYE Packed Column


Column Size I.D. x Length (mm)	Product Number
4.6 x 250	37989-11
10 x 250	37996-11
20 x 250	38044-41
28 x 250	34300-91

COSMOSIL 5PYE Guard Column

Column Size I.D. x Length (mm)	Product Number
4.6 x 10	37903-11
10 x 20	38041-71
20 x 50	34475-21

Fullerene Chromatogram Index

Fullerene Chromatogram Index includes more than 100 chromatograms. If you are interested in this index, please feel free to e-mail us at info.intl@nacalai.com. The online version is available at the website of The Fullerenes, Nanotubes and Graphene Research Society below.



The Fullerenes, Nanotubes and Graphene Research Society

Website: http://fullerene-jp.org/en/chromato_index_3.pdf

(7) Columns for Soluble Carbon Nanotube Separation

COSMOSIL CNT-300, CNT-1000, CNT-2000

- Size-based separation of soluble carbon nanotubes
- Three pore sizes (300 Å , 1000 Å , 2000 Å)
- High durability


Specifications

Packing Material	CNT-300	CNT-1000	CNT-2000
Silica Gel	High purity porous spherical silica		
Average Particle Size	5 µm		
Average Pore Size	approx. 300 Å	approx. 1000 Å	approx. 2000 Å
Bonded Phase	Hydrophilic group (neutral)		
pH Range	2-7.5		
Pressure	15 MPa and below		

Applications

• Carbon Nanotubes

COSMOSIL CNT columns offer improved separation for DNA-wrapped carbon nanotubes by connecting three columns with different pore sizes.



Ordering Information

- Analytical Columns (Particle Size: 5µm)

COSMOSIL CNT-300 Packed Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 300	09195-71

COSMOSIL CNT-300 Guard Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 50	09194-81

COSMOSIL CNT-1000 Packed Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 300	09197-51

COSMOSIL CNT-1000 Guard Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 50	09196-61

COSMOSIL CNT-2000 Packed Column

Column Size I.D. x Length (mm)	Product Number
7.5 x 300	09199-31

COSMOSIL CNT-2000 Guard Column






Column Size I.D. x Length (mm)	Product Number
7.5 x 50	09198-41

II. SFC Columns

Supercritical Fluid Chromatography (SFC) has become more attractive because it offers some advantages over HPLC, such as high speed, unique selectivity and environmentally friendly separations. Many conventional normal-phase stationary phases, such as diol, amino and cyano, have been used for SFC applications. However, these phases present limitations for separations. COSMOSIL SFC Columns have been developed to enhance the capability of SFC separations.

COSMOSIL SFC Columns

Nacalai Tesque has developed columns specially designed for SFC in collaboration with Nacalai USA and Pfizer, Inc. Global R&D: COSMOSIL HP, PY (equivalent to 2-ethylpyridine) and Quinoline. In addition to these, our HPLC columns Cholester and PBr have been tested for use with SFC.



Packing Material	HP	PY	Quinoline		Cholester		PBr
Average Particle Sizes	3, 5 μm		2.5 μm	5 μm	2.5 μm	5 μm	5 μm
Average Pore Sizes	120 $\text{\AA}$		130 $\text{\AA}$	120 $\text{\AA}$	130 $\text{\AA}$	120 $\text{\AA}$	120 $\text{\AA}$
Specific Surface Area	300 m^2/g		330 m^2/g	300 m^2/g	330 m^2/g	300 m^2/g	300 m^2/g
Bonded Phase Structure							
Bonded Phase	3-Hydroxyphenyl group	Pyridinyl group	Quinoline group		Cholesteryl group		Pentabromobenzyl group

Pharmaceutical Analysis

Each phase has different retention properties.

• Imipramine



Column: COSMOSIL **
 Column size: 2.1mm I.D.-150mm
 Mobile phase:
 A: CO_2
 B: 0.1% $\text{CH}_3\text{COONH}_4$ - Methanol
 B conc. 0→60% (0→14min), 60% (14-17min)
 Flow rate: 0.8 ml/min
 BPR: 10 MPa
 Temperature: 40 $^\circ\text{C}$
 Detection: UV260nm
 Sample: Imipramine (1mmol/l)
 Inj. Vol.: 2.0 μl



Data courtesy of Kyushu University Medical Institute of Bioregulation Research Center for Transomics Medicine Division of Metabolomics SFC-214

• Procaine


Column: COSMOSIL **
 Column size: 2.1mm I.D.-150mm
 Mobile phase:
 A: CO_2
 B: 0.1% $\text{CH}_3\text{COONH}_4$ - Methanol
 B conc. 0→60% (0→14min), 60% (14-17min)
 Flow rate: 0.8 ml/min
 BPR: 10 MPa
 Temperature: 40 $^\circ\text{C}$
 Detection: UV280nm
 Sample: Procaine (1mmol/l)
 Inj. Vol.: 2.0 μl



Data courtesy of Kyushu University Medical Institute of Bioregulation Research Center for Transomics Medicine Division of Metabolomics SFC-210

Comparison of Retention Behavior


The following three stationary phases were evaluated for their retention of hydrophilic, hydrophobic, acidic and basic compounds. COSMOSIL HP and PY elute hydrophobic compounds first and retain hydrophilic compounds longer, whereas PBr elutes in the reverse order, exhibiting high retention for hydrophobic compounds. HP had the longest retention for basic compounds.




COSMOSIL HP (3-Hydroxyphenyl)

Applications

• Non-steroidal anti-inflammatory drugs



• Beta Blockers (Peak elution order reversal under identical conditions)



Ordering Information

- Analytical / Preparative Columns

(Particle Size: 5 µm)

COSMOSIL 5HP (3-Hydroxyphenyl) Packed Columns

Column Size I.D. x Length (mm)	Product Number
2.0 x 150	13787-91
4.6 x 250	13788-81
10 x 250	13789-71
20 x 250	13790-31

COSMOSIL 5HP (3-Hydroxyphenyl) Guard Columns

Column Size I.D. x Length (mm)	Product Number
10.0 x 20	13791-21

(Particle Size: 3 µm)


COSMOSIL 3HP (3-Hydroxyphenyl) Packed Columns

Column Size I.D. x Length (mm)	Product Number
2.0 x 150	13792-11
4.6 x 250	13793-01

COSMOSIL PY (Pyridinyl)

Applications

- Hydrophilic organics



Ordering Information

- Analytical / Preparative Columns

(Particle Size: 5 µm)

COSMOSIL 5PY (Pyridinyl) Packed Columns

Column Size I.D. x Length (mm)	Product Number
2.0 x 150	13818-81
4.6 x 250	13827-61
10 x 250	13828-51
20 x 250	13829-41

COSMOSIL 5PY (Pyridinyl) Guard Columns

Column Size I.D. x Length (mm)	Product Number
10.0 x 20	13830-01

(Particle Size: 3 µm)

COSMOSIL 3PY (Pyridinyl) Packed Columns


Column Size I.D. x Length (mm)	Product Number
2.0 x 150	13831-91
4.6 x 250	13832-81

COSMOSIL Quinoline


The structural similarities between polar lipids, such as cholesterol and related analogs, have posed chromatographic and spectrometric challenges to analysts interested in quantifying these potential biomarkers. COSMOSIL Quinoline has been developed to improve the separation of these structural isomers utilizing the π - π interactions and structural rigidity of the naphthylethyl phase and the hydrogen bonding of the pyridine phase.

Applications

• Steroids



• Caffeine analogs



Ordering Information

• Analytical / Preparative Columns

(Particle Size: 5 μ m)

COSMOSIL Quinoline Packed Column

Column Size I.D. x Length (mm)	Product Number
2.0 x 150	Inquire
4.6 x 100	Inquire
4.6 x 150	Inquire
10.0 x 150	Inquire
20.0 x 150	Inquire

(Particle Size: 2.5 μ m)


COSMOSIL Quinoline Packed Column

Column Size I.D. x Length (mm)	Product Number
3.0 x 50	Inquire
3.0 x 100	Inquire
3.0 x 150	Inquire

COSMOSIL Cholester

Fat-Soluble Vitamin Analysis

When used with SFC, COSMOSIL Cholester can separate fat-soluble vitamins and their impurities.



COSMOSIL Cholester exhibits strong retention for fat-soluble vitamins and is suitable for on-line SFE-SFC using Shimadzu's Nexera UC. The online extraction from food also produced triglyceride impurities, which were successfully separated from the vitamins.

Ordering Information

For ordering information for COSMOSIL Cholester, refer to page 24.

COSMOSIL PBr

Ordering Information

For ordering information for COSMOSIL PBr, refer to page 25.

III. Preparative Packing Materials

Normal and Reversed Phase Packing Materials


Introduction

Open column chromatography is an excellent and easy technique for large-scale preparation and purification at low cost. COSMOSIL offers both normal and reversed phase packing materials based on totally porous spherical silica, which provides higher separation, less pressure and higher reproducibility than irregular silica.

Specifications

Packing Material	C ₁₈ -OPN	C ₁₈ -PREP	Silica Gel 60 (Neutral)
Silica Gel	High purity porous spherical silica		
Average Particle Size	75, 140 µm	50, 75, 140 µm	75, 140 µm
Average Pore Size	approx. 120 Å		approx. 60 Å
Specific Surface Area	approx. 300 m ² /g		approx. 500 m ² /g
Bonded Phase	Octadecyl group		None
Carbon Content	—	approx. 19%	0%
Residual Silanol Group	Yes	None	—
Application	Open column chromatography / Flash column chromatography		
	Reversed phase chromatography		Normal phase chromatography

Selection Guide (Reversed Phase)



COSMOSIL C₁₈-OPN

- A new “Water-Wet” C₁₈ packing material for reversed phase open column chromatography
- Usable under 100% aqueous eluents

Characteristic

The external surface of the C₁₈-OPN gel is coated with hydrophilic group to increase wettability of the gel, and the octadecyl group is bonded within the pore of the gel. Conventional reversed phase C₁₈ packing materials are restricted to about 30-50% water in the mobile phase. The COSMOSIL C₁₈-OPN is a new “Water-Wet” C₁₈ packing material developed for reversed phase open column chromatography. The C₁₈-OPN material can be used in 100% aqueous eluents.




Figure 1. Structure of C₁₈-OPN



Figure 2. Packing material in water

Left: C₁₈-OPN provides good resolution


Can be used with low concentration of organic solvent on open, flash column chromatography.

Right: C₁₈-PREP float up

Use with 70% or more organic solvent on open, flash column chromatography.

Applications

- Separation of hydrophilic compounds in aqueous solution



In reversed phase chromatography, hydrophilic compounds such as theobromine and theophylline could be separated under low concentration of organic solvent. The figure shows they are clearly separated by reversed open column chromatography with 70% water.

Ordering Information

COSMOSIL C₁₈-OPN

Product Name	Average Particle Size	Product Number	PKG Size
COSMOSIL 75C ₁₈ -OPN	75μm	37842-66	100 g
		37842-95	500 g
		37842-11	1 kg
COSMOSIL 140C ₁₈ -OPN	140μm	37878-16	100 g
		37878-45	500 g
		37878-61	1 kg

COSMOSIL C₁₈-PREP

- Standard reversed phase packing material for open chromatography
- Endcapped
- 3 particle sizes (50, 75, 140 μm)

Particle Size, Flow Rate and Theoretical Plate Number

Because reversed phase chromatography employs a mobile phase of high viscosity such as methanol and water, the flow rate is lower than that of normal phase chromatography, which uses mobile phase of low viscosity such as hexane and ethyl acetate.




Figure 1. Flow rate against theoretical plate number
Column size: 20 mm I.D. x 300 mm





Figure 2. Concentration of methanol against flow rate
Column size: 10 mm I.D. x 180 mm bed height
(gravitational liquid flow)

Performance Evaluation


COSMOSIL Application Data

Column: 50C₁₈-PREP
Column size: 4.6mm I.D.-250mm (Closed column)
Mobile phase: Methanol/Water = 60 / 40
Temperature: 30°C
Detection: UV 254 nm
Flow rate: 2 ml/min
Sample: (Methanol solution)
1; Uracil (0.15 mg/mL)
2; Acetophenone (0.2 mg/mL)
3; Benzene (20 mg/mL)
4; Toluene (40 mg/mL)
Inj. Vol: 5 μL



COSMOSIL Application Data

Column: 75C₁₈-PREP
Column size: 4.6mm I.D.-250mm (Closed column)
Mobile phase: Methanol/Water = 30 / 70
Temperature: 30°C
Detection: UV 254 nm
Flow rate: 1 ml/min
Sample: (Methanol solution)
1; Uracil (0.04 mg/mL)
2; Caffeine (0.5 mg/mL)
3; Phenol (2 mg/mL)
Inj. Vol: 20 μL



Ordering Information

COSMOSIL C₁₈-PREP


Product Name	Average Particle Size	Product Number	PKG Size
COSMOSIL 50C ₁₈ -PREP	50 μm	12065-84	100 g
		12065-55	500 g
		12065-71	1 kg
COSMOSIL 75C ₁₈ -PREP	75 μm	12061-24	100 g
		12061-95	500 g
		12061-11	1 kg
COSMOSIL 140C ₁₈ -PREP	140 μm	12063-04	100 g
		12063-75	500 g
		12063-91	1 kg

Silica Gel (Spherical, Neutral)


- The pH of silica gel is adjusted to neutral
- Suitable for the separation of pH-sensitive compounds

Comparison with Conventional Silica Gel

• Purification of Acetals -1



• Purification of Acetals -2



Ordering Information

Silica gel 60 (Spherical, Neutral)

Product Name	Average Particle Size	Product Number	PKG Size
Silica Gel 60 (Spherical, Neutral) for Column Chromatograph	75 µm	30511-64	100 g
		30511-35	500 g
		30511-51	1 kg
		30511-06	5 kg
		30511-22	25 kg
	140 µm	30518-94	100 g
		30518-65	500 g
		30518-81	1 kg

Silica Gel (for Column Chromatography)

[Ordering Information](#)

Silica Gel (Spherical)

Product Name	Average Particle Size	Average Pore Size	Grade	Product Number	PKG Size
Silica Gel 60, Spherical	approx. 70 ~ 230 mesh	60 Å	SP	30731-71	1 kg
				30731-42	25 kg
Silica Gel 120, Spherical	approx. 70 ~ 230 mesh	120 Å	SP	30734-41	1 kg

Silica Gel (Irregular)

Product Name	Average Particle Size	Average Pore Size	Grade	Product Number	PKG Size
Silica Gel 60	approx. 70 ~ 230 mesh	60 Å	SP	30724-55	500 g
				30724-71	1 kg
				30724-84	5 kg
				30724-42	25 kg
	approx. 230 ~ 400 mesh	60 Å	SP	30721-85	500 g
				30721-01	1 kg
				30721-14	5 kg

IV. Related Products


1. Reagents for Mobile Phase Preparation

Phosphate Buffer Solution (pH 2.5) (5X)

- pH-adjusted
- UV, fluorescence tested
- Filtered (0.2 µm)
- Easily prepare the mobile phases used in COSMOSIL applications

How to Prepare

Dilute this product with HPLC grade distilled water (1 part buffer solution : 4 parts water) to make the 20 mmol/l phosphate buffer used in the following COSMOSIL applications.



Ordering Information

Phosphate Buffer Solution (5X)

Product Name	Grade	Product Number	PKG Size
Phosphate Buffer Solution (pH 2.5) (5X)	SP	08969-71	1 L
Phosphate Buffer Solution (pH 7.0) (5X)	SP	08968-81	1 L

Stock Solutions for HPLC

Ordering Information

Product Name	Grade	Product Number	PKG Size
1mol/l-Ammonium Formate Solution	SP	12235-54	100 ml
1mol/l-Ammonium Acetate Solution	SP	12236-44	100 ml

Premixed Eluents for HPLC

Ordering Information

Product Name	Grade	Product Number	PKG Size
0.1vol% Formic Acid-Acetonitrile	SP	12578-61	1 L
		12578-03	3 L
0.1vol% Formic Acid-Distilled Water	SP	12582-91	1 L
		12582-33	3 L
0.1vol% Trifluoroacetic Acid-Acetonitrile	SP	12583-81	1 L
		12583-23	3 L
0.1vol% Trifluoroacetic Acid-Distilled Water	SP	12584-13	3 L

Additives

Ordering Information

Product Name	Grade	Product Number	PKG Size
Acetic Acid	SP	08963-02	25 ml
Formic Acid	SP	08965-82	25 ml
Phosphoric Acid, Ortho	SP	08964-92	25 ml
Trifluoroacetic Acid	SP	34840-21	5 x 1 ml
		34840-76	5 x 1.5 ml
		34840-63	5 x 3 ml
		34840-34	10 ml

Ion-pair Reagents

Ordering Information

For Basic Samples

(R-SO₃⁻Na⁺)

Product Name	R:	Grade	Product Number	PKG Size
Sodium 1-Butanesulfonate	C ₄ H ₉ -	SP	31331-94	5 g
Sodium 1-Pentanesulfonate	C ₅ H ₁₁ -	SP	31730-64	5 g
			31730-22	25 g
Sodium 1-Hexanesulfonate	C ₆ H ₁₃ -	SP	31529-24	5 g
			31529-82	25 g
Sodium 1-Heptanesulfonate	C ₇ H ₁₅ -	SP	31528-34	5 g
			31528-92	25 g
Sodium 1-Octanesulfonate	C ₈ H ₁₇ -	SP	31729-04	5 g
			31729-62	25 g
Sodium 1-Nonanesulfonate	C ₉ H ₁₉ -	SP	31626-44	5 g
Sodium 1-Decanesulfonate	C ₁₀ H ₂₁ -	SP	31429-34	5 g
Sodium 1-Dodecanesulfonate	C ₁₂ H ₂₅ -	SP	31426-64	5 g
Sodium Lauryl Sulfate	**	SP	31623-32	25 g

0.5M Solution

Sodium 1-Butanesulfonate	C ₄ H ₉ -	SP	31332-84	5 x 10 ml
Sodium 1-Hexanesulfonate	C ₆ H ₁₃ -	SP	31532-64	10 ml
			31532-06	5 x 10 ml
Sodium 1-Octanesulfonate	C ₈ H ₁₇ -	SP	31733-34	10 ml
			31733-76	5 x 10 ml

For Acidic Samples

(C₄H₉)₄N⁺X⁻

Product Name	X ⁻ :	Grade	Product Number	PKG Size
Tetra- <i>n</i> -butylammonium Bromide	-Br	SP	32824-72	25 g
Tetra- <i>n</i> -butylammonium Chloride	-Cl	EP	32935-64	5 g
			32935-22	25 g
Tetra- <i>n</i> -butylammonium Hydrogensulfate	-HSO ₄	GR	32924-62	25 g
Tetra- <i>n</i> -butylammonium Iodide	-I	SP	32905-54	5 g
			32905-12	25 g
Tetra- <i>n</i> -butylammonium Perchlorate	-ClO ₄	SP	32906-44	5 g
			32906-02	25 g
Tetra- <i>n</i> -butylammonium Phosphate	-H ₂ PO ₄	SP	32929-54	5 g

0.5M Solution

Tetra- <i>n</i> -butylammonium Phosphate	-H ₂ PO ₄	SP	32926-26	10 ml
			32926-84	5 x 10 ml

2. Products for Sample Preparation

Cosmonice Filter


- For sample filtration
- Just attach a filter on top of a syringe

W Series (aqueous solution)

The W series uses a new material of low-adsorptive and low-extractive PVDF (poly vinylidenedifluoride) filter, which can be used with various solvents. They are able to minimize the loss of proteins in the small amount of sample, and prevent secondary contamination during prefiltration.

S Series (organic solvents)

The S series uses a PTFE (poly tetrafluoroethylene) filter, which shows strong resistance for solvents, acids, and alkalis. It is best for prefiltration of samples extracted with solvents such as chloroform and tetrahydrofuran.



Ordering Information

Cosmonice Filter

Product Name	Diameter (mm)	Pore Size (μm)	Process Volume	Hold-up Volume	Product Number	PKG Size
Cosmonice Filter W (aqueous)	4	0.45	1 ml or less	< 10 μl	06543-04	100 pkg
	13	0.45	0.5~10 ml	< 30 μl	06544-94	100 pkg
Cosmonice Filter S (solvent)	4	0.45	1 ml or less	< 10 μl	06541-24	100 pkg
	13	0.45	0.5~10 ml	< 30 μl	06542-14	100 pkg

Connection Inlet: luer-lock; Outlet: luer-slip, Connectable to needles
Housing : polyethylene

Cosmospin Filter

- For sample filtration
- Easy to use by centrifugation
- Omnipore hydrophilic PTFE membrane filter

Ordering Information

Cosmospin Filter

Product Name	Pore Size (μm)	Maximum Sample Volume	Hold-up Volume	Maximum Centrifugal Force	Rotor Size (fixed-angle)	Filtration Area	Color	Product Number	PKG Size
Cosmospin Filter G	0.2	0.4 ml	5 μl	5000 x g	1.5 ml	0.2 cm ²	Brown	06549-44	100 pkg
Cosmospin Filter H	0.45	0.4 ml	5 μl	5000 x g	1.5 ml	0.2 cm ²	White	06540-34	100 pkg

Dimensions: 10.6 mm diameter x 45 mm
Membrane: Omnipore hydrophilic PTFE
Sample reservoir and collection tube: Polypropylene



Labeling Reagents

Ordering Information

Product Name	Grade	Storage	Product Number	PKG Size
Dabsyl Chloride	SP	Room temp.	10427-91	1 g
3,5-Dinitrobenzoyl Chloride (DNBC)	SP	Dark and Cool	13530-44	5 g
NBD Chloride	SP	Refrigerator	24113-61	1 g
o-Phthalaldehyde (OPA)	SP	Refrigerator	27824-61	1 g
			27824-74	5 g
			27824-32	25 g

I. HPLC Columns

II. SFC Columns

III. Preparative Packing Materials

IV. Related Products

3. Column Care Products

Introduction

It is important to preserve a column by washing it with suitable cleaning methods before storing it under appropriate conditions to obtain stable data and prolong the column lifetime.

Applicable Columns

Cleaning Solution Kit and Storage Solution for Reversed Phase HPLC Columns is only applicable to reversed phase HPLC columns, such as COSMOSIL C₁₈-MS-II, AR-II, PAQ, EB, Cholester, πNAP, PYE, PBr and COSMOCORE 2.6C₁₈, 2.6Cholester and 2.6PBr. Please note that this product is not suitable for Sugar-D, HILIC, normal phase or ion exchange columns.


Cleaning Solution Kit for Reversed Phase HPLC Columns

Components

Product Name	Main Components	PKG Size	Quantity	Container
Cleaning Solution A	Methanol	500 ml	2	Brown Glass Bottle
Cleaning Solution B	Tetrahydrofuran, Methanol	500 ml	1	Brown Glass Bottle

Application

Cleaning Solution Kit for Reversed Phase HPLC Columns is designed for washing away contaminant adsorption and stationary phase shedding. If you experience the following symptoms, please try their corresponding solution first.




Procedure

(For 4.6 mm I.D. x 150 mm)

- Replace solvent with HPLC-grade distilled water (1 ml/min, 30 min).
(*This step is for mobile phases containing high concentration buffer. If you are using a salt-free mobile phase, please start from step (2).)
- Run the Cleaning Solution A through the column for 15 min at a flow rate of 1ml/min.
- Run the Cleaning Solution B through the column at a flow rate of 1ml/min until the baseline becomes stable (approx. 15 min).
- Run the Cleaning Solution A through the column for 15 min. The column is ready for storage.

Example of pressure difference between washed and unwashed columns

The figure shows a pressure comparison between washed and unwashed columns using Cleaning Solution Kit. Repeated analysis of natural products was conducted using COSMOSIL 5C₁₈-MS-II (4.6 mm I.D. x 150 mm).



(Condition)

Column: COSMOSIL 5C₁₈-MS-II (4.6 mm I.D. x 150 mm)

Mobile phase: Methanol / H₂O = 70 / 30


Flow rate: 1.0ml/min

Temperature: 40°C

As shown in the figure above, the column pressure increases if you use column continuously without washing. If you wash the column, you can extend the column life time and ease the pressure burden on your HPLC equipment.

Example of a Stable Baseline

The baseline may be unstable if sample components with very long retention remain in the column or stationary phase shedding occurs. Especially when analyzing crude samples that have components with a wide range of chemical characteristics, some unwanted components may be strongly retained in the column and slowly elute out in subsequent runs. The resulting unstable baseline can be eliminated by washing the column with the Cleaning Solution Kit.



Ordering Information

Product Name	Grade	Product Number	PKG Size
Cleaning Solution for Reversed Phase HPLC Columns	SP	08966-30	1 kit

Storage Solution for Reversed Phase HPLC Columns

Storage Solution for Reversed Phase HPLC Columns is designed for storing columns under suitable conditions.

Procedure

(For 4.6 mm I.D. x 150 mm)

(1) Replace solvent with HPLC-grade distilled water. (1 ml/min, 30 min)

(*This step is for mobile phases containing high concentration buffer. If you are using a salt-free mobile phase, please start from step(2).)

(2) Run the Storage Solution through the column for 15 min at a flow rate of 1ml/min, and store.

Ordering Information

Product Name	Grade	Product Number	PKG Size
Storage Solution for Reversed Phase HPLC Columns	SP	08967-20	1 kit (500 ml)

4. COSMOSIL HPLC Accessories

Ordering Information

COSMOSIL Guard Cartridge Holder

I.D.	Product Number	PKG Size
2.0 mm	11884-71	1 PKG
4.6 mm	38009-79	1 PKG



Guard Cartridge Holder is required for Guard Cartridge.

COSMOSIL Column Prefilter

Product Number	PKG Size
39361-19	1 PKG



COSMOSIL Column Prefilter employs filter with smaller pore size (1 μm) than that of column frit (2 μm) .

COSMOSIL Column Spare Filter for Prefilter

Product Number	PKG Size
39539-09	2 PKG



Column spare filter for prefilter

COSMOSIL Column Connecting Tube

I.D.	Product Number	PKG Size
0.1 mm	12570-41	1 PKG
0.25 mm	37843-69	1 PKG



For connecting columns



COSMOSIL

COSMOSIL Technical Notes

For our COSMOSIL FAQ, troubleshooting, and technical information, please visit our web site at <http://www.nacalai.co.jp/global/cosmosil/>.

COSMOSIL HPLC Columns

General Info. of COSMOSIL/COSMOGEL
COSMOSIL Columns List by Phase

- Standard Reversed Phase Columns
- Specialty Reversed Phase Columns
- Ultra-High Performance Columns
- Normal Phase Columns

Related Products

- Preparative Packing Materials
- Related Products
- Prefiltration Tools
- Fatty Acid Methylation **NEW**

► Natural Compounds
Crude Drug Standards
Plant Extract Standards

- Hydrophilic Interaction Columns
- Saccharide Separation Columns
- Protein Separation Columns (Wide Pore Columns)
- Fullerene Separation Columns
- Carbon Nanotubes Separation Columns

► COSMOSIL Applications
Application Search
Over 7,000 Data

► Reference Lists
Reference Lists

► Technical Notes
FAQ
Troubleshooting
Technical Information

Click

Technical Notes

FAQ and Troubleshooting

[FAQ and Troubleshooting](#)

Click

Technical Information

1. Preparation of Mobile Phase for HPLC (PDF 158 KB)
2. Inner Diameter of Column(scale down and scale up) (PDF 344 KB)
3. Troubleshooting for Increased Pressure (PDF 149 KB)
4. Sample Pretreatment for HPLC (PDF 682 KB)
5. Baseline Noise in Gradient Elution (PDF 155 KB)
6. Effect of Guard Column (PDF 642 KB)
7. Selectivity of Packing Materials in Reversed Phase Liquid Chromatography (PDF 1,504 KB)
8. Methods in Developing Mobile Phase Condition for C18 Column (PDF 253 KB)
9. Comparison with Old Type COSMOSIL (PDF 473 KB)

Click

FAQ and Troubleshooting

• FAQ

Q1	What is the pressure limit of column?
Q2	What is the flow rate limit?
Q3	What is the recommended pH range?
Q4	What is the concentration of buffer and salt?
Q5	How do I adjust mobile phase?
Q6	What solvent grade should I use for the mobile phase?
Q7	What is the difference between acetonitrile and methanol?
Q8	Which mobile phase can be used for LC/MS or ELSD detector?
Q9	What should I pay attention to when I use ion-pairing reagents?
Q10	What flow direction should I use for the mobile phase?
Q11	What is the recommended temperature range of columns?
Q12	What is the shipping solvent?
Q13	How do I wash columns?
Q14	How do I store columns?
Q15	How long does a column last?

3. Troubleshooting for Increased Pressure


Introduction

Repeated analysis may increase back pressure. Continuous use of HPLC columns under high pressure can cause deterioration and overload of the equipment. Therefore, it is important to monitor column back pressure regularly and solve the problem timely.

Identification of the Clogging Site

The back pressure increase can be due to clogging of a column or clogging of the equipment. First of all, identify the clogging site.

Schematic Diagram of HPLC





Follow us!
@NacalaiTesque

Stay up to date and get useful tips!



Warranties and Disclaimers:

Nacalai Tesque warrants that its products shall conform to the description of such products as provided by Nacalai Tesque through its catalog, analytical data or other literature. Nacalai Tesque makes no other warranty, express or implied, as to the fitness of these products for any particular purpose. Nacalai Tesque shall not in any event be liable for incidental or consequential damages that may result from any use or failure of the products.

For more information on products and pricing, please contact your local distributor.

NACALAI TESQUE, INC.

Nijo Karasuma, Nakagyo-ku, Kyoto
604-0855 JAPAN

TEL : +81 (0)75 251 1730

FAX : +81 (0)75 251 1763

E-mail : info.intl@nacalai.com

www.nacalai.com

Nacalai USA, Inc.

10225 Barnes Canyon Rd., Suite A 103
San Diego, CA 92121

TEL : (858) 404-0403

FAX : (858) 404-0408

E-mail : info@nacalaiusa.com

www.nacalaiusa.com